

# **Rate Constant and Temperature Dependence for the Reaction of Hydroxyl Radicals with 2-Fluoropropane (FC-281ea) and Comparison with an Estimated Rate Constant**

**W. B. DeMore and Edmond W. Wilson, Jr.**

*Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109*

## **Abstract**

Relative rate experiments were used to measure the rate constant and temperature dependence of the reaction of OH radicals with 2-fluoropropane (HFC-281ea), using ethane, propane and ethyl chloride as reference standards. Measurements were made using both infrared spectroscopy and gas chromatography (GC). Results from the two measurement techniques were in good agreement, with the GC data being more precise. The rate at 298 K for HFC-281ea, based on the GC experiments, was found to be  $5.7\text{E-}13 \text{ cm}^3/\text{molec.-s.}$ , with Arrhenius A-factor =  $3.06\text{E-}12 \text{ cm}^3/\text{molec.-s.}$  and  $E/R = 503 \text{ K}$ . A previously described estimation technique, along with an improved method to estimate the Arrhenius parameters, is used to predict a rate constant and temperature dependence for 2-fluoropropane which is in good agreement with the experimental value. A prediction is also made for 1-fluoropropane, for which there are no data.

## **Introduction**

It is increasingly evident that accurate assessment must be made of the consequences of injection of gaseous materials into the atmosphere that arise from urban, industrial and natural sources. For molecules containing at least one hydrogen atom, reaction with tropospheric hydroxyl radicals is the primary pathway of removal of these compounds from the atmosphere. Therefore, atmospheric lifetimes of these gases are determined mainly by the rate at which they react with tropospheric hydroxyl radicals. Predictive rules that can accurately estimate

atmospheric lifetimes are useful, and we are measuring the kinetics of key compounds to use as a basis set for forming predictive rules. In this connection, the rate constant and temperature dependence for the reaction of 2-fluoropropane, HFC-281ea, with hydroxyl radical are reported in this article. We shall examine the consequences of this information on empirical rules previously developed to predict rate constants and temperature dependencies.<sup>2</sup>

## Experimental

The technique used in this work has been described in several recent publications.<sup>2,6,5</sup> The reaction measured is



A relative method is used in which the fractional loss of reactants is compared to the fractional loss of a reference compound. The fractional loss is reported in terms of a depletion factor, DF, which is the ratio of initial concentration to final concentration. The experiments were carried out in either a stopped-flow mode or a stirred-flow mode. For the stopped-flow mode the relation of sample to reference rate constant is given by

$$\frac{k_{\text{reactant}}}{k_{\text{reference}}} = \frac{\ln(DF)_{\text{reactant}}}{\ln(DF)_{\text{reference}}} \quad (2)$$

For the stirred-flow mode the rate constant ratio is given by

$$\frac{k_{\text{reactant}}}{k_{\text{reference}}} = \frac{[(DF)_{\text{reactant}} - 1]}{[(DF)_{\text{reference}} - 1]} \quad (3)$$

The OH radicals were produced by direct photolysis of H<sub>2</sub>O at 185 nm or by 254 nm photolysis of O<sub>3</sub> (5-10 x 10<sup>16</sup> cm<sup>-3</sup>) in the presence of water vapor (3-5 x 10<sup>17</sup> cm<sup>-3</sup>). The experiments were done in temperature-controlled cylindrical quartz cells. The cylindrical cells were 10 cm in length and 5 cm in diameter and water-jacketed for measurements below 95°C or wrapped with heating tape and insulating material for higher temperatures. Residence times in the cells for the stirred-flow mode were about one minute.

All concentration measurements were made either by gas chromatography or infrared spectroscopy. For the gas chromatography measurements an SRI Model 8610 gas chromatograph equipped with flame ionization detector was used. Samples were injected into the instrument by means of a thermostated 1.00 mL sample loop. The sample loop was evacuated and the sample allowed to flow into the loop. An electronically operated valve sent the sample into the He carrier gas stream of the chromatograph. The PeakSimple Data System, Version 1.59, for Windows 3.1 controlled instrument operation, data collection and peak integrations. A 6 ft., 1/8 in. o.d. Poropak P, 80/100 mesh column, operated at 60°C or 80°C was used for most of the separations. A 3 ft., 1/8 in. o.d. silica gel, 80/100 mesh column, operated at 175°C was used for reactions involving chloroethane. Infrared measurements were taken with a Nicolet 20SX FTIR, operated at 0.5 cm<sup>-1</sup> resolution in the absorbance mode. Samples were contained in a White cell of 3 m. pathlength. In order to minimize the risk of analytical errors, such as overlapping peaks in the GC analysis, measurements were made using both infrared spectroscopy and gas chromatography to measure the depletion factors as a function of temperature. Although both techniques give essentially the same results, the data obtained by the gas chromatography method have significantly less random error, mainly because the IR method is less precise for hydrocarbons, in which the C-H bands must be relied on for the analysis.

## Results

**Table 1** gives the ratio results using infrared spectroscopy for HFC-281ea vs three reference compounds; ethane, propane and chloroethane. These measurements were made in a stirred-flow mode using Equation 3 to calculate the rate ratios. **Table 2** presents the ratio results using gas chromatography for ethane and propane as reference compounds. Stopped-flow methods were utilized in the gas chromatography measurements, and rate constant ratios were calculated from Equation 2. The absolute rates for the reaction of HFC-281ea with hydroxyl radical were calculated using both IR and GC data. The data sets are in agreement, but the GC

data are preferable due to the lower random error. **Table 3** gives the values of the rates of the reference materials used to calculate the rate and temperature dependence for HFC-281ea.

The resulting rate constant expression for HFC-281ea, based on the GC ratios vs ethane and propane, is

$$k(281ea) = 3.06E-12 \exp(-503/T) \text{ cm}^3/\text{molec.-s.} \quad (4)$$

$$k(298 \text{ K}) = 5.7E-13 \text{ cm}^3/\text{molec.-s.} \quad (5)$$

The uncertainty in  $k(298 \text{ K})$  is about 10% and the uncertainty in  $E/R$  is about 100 K. **Figure 1** shows an Arrhenius plot of the data along with the least squares fit corresponding to Equation 4. **Figure 2** compares the IR data with Equation 4, showing that although these data are more scattered, they are in good agreement with the GC data.

## Discussion

It is useful to compare the measured rate expression for 281ea with an estimated rate expression based on a method previously described.<sup>2</sup> According to this method, each reactive site is considered to be a methane derivative, and the reactivity of the C-H bonds at that site is determined by the attached groups which have replaced the H-atoms in methane. The total rate constant is obtained as the sum of the contributions from each C-H bond present in the molecule. The logarithm (base 10) of the rate constant (per H-atom) at each site is given by:

$$\log k = \log k(\text{CH}_4) + G_1 \cdots G_3 \quad (6)$$

The quantity  $\log k(\text{CH}_4)$  is the logarithm of the  $\text{CH}_4$  rate constant per H-atom, and is equal to -14.79, corresponding to  $k(\text{CH}_4) = 6.5E-15 \text{ cm}^3/\text{molec.-s.}$  at 298 K. The G-terms are the contributions for the various groups, such as Cl, Br,  $\text{CH}_3$ ,  $\text{CF}_3$ , etc., which may be present on the carbon atom in place of an H-atom. The G-terms are determined by a fitting procedure based on a large number of well-determined rate constants at 298 K.<sup>2</sup> Thus the use of Equation 6 yields an estimate of the rate constant at 298 K. The current set of G-values is listed in **Table 4**.

To determine the  $E/R$  or activation temperature of the reaction, we use an estimated A-factor based on a database of reactions similar to that used to calibrate the G-values at 298 K. In

our previous description of the estimation technique, we used a fixed A-factor of  $8\text{E-}13$   $\text{cm}^3/\text{molec.-s.}$  per H-atom. However, the database now includes reactions over a wider range of rate constants, and a clear dependence of A-factor on reaction rate is now apparent. The database for this calibration is listed in **Table 5**, which includes those A-factors from our previous work which are in good agreement with independent measurements by absolute methods. A plot of A-factor per H atom vs  $\log(k/n(\text{H}))$  for rate constants ranging over four orders of magnitude is shown in **Figure 3**. The scatter in this plot is similar in magnitude to the uncertainty in the measured A-factors, but may well be due in part to real variations in the A-factors for these reactions. The least squares fit to these data is as follows:

$$\log A = 0.17 \log(k/n) - 9.61 \quad (7)$$

The procedure therefore is to estimate the 298 K rate constant per H-atom at each reactive site using Equation 6, and then to calculate the A-factor for these C-H bonds using Equation 7. The E/R value is then calculated from the expression,

$$E/R = -298 \ln(k/A) \quad (8)$$

The total reaction rate at a given temperature is obtained by summing the contributions from each site. The effective overall Arrhenius parameters for the reaction are then obtained by fitting the sums over the desired temperature range to a single Arrhenius equation. This procedure is illustrated below for both n-propyl fluoride and isopropyl fluoride, which serves to demonstrate the interplay between A-factors and activation temperatures factors for these two reactions.

*Estimated Rate Constant for  $\text{CH}_3\text{CH}_2\text{CH}_2\text{F}$  :*

$$-\text{CH}_3 \text{ site: } \log k/n(\text{H}) = -14.79 + G(\text{CH}_3) + \log 3 = -13.54$$

(The  $\text{CH}_3$  group has been used as a surrogate for the  $\text{CH}_2\text{CH}_2\text{F}$  group.)

$$k(298 \text{ K}) = 8.7\text{E-}14 \text{ cm}^3/\text{molec.-s.}$$

$$A = 3.7\text{E-}12 \text{ cm}^3/\text{molec.-s.}$$

$$E/R = 1117 \text{ K}$$

$$-\text{CH}_2- \text{ site: } \log k/n(\text{H}) = -14.79 + G(\text{CH}_3) + G(\text{CH}_2\text{F}) = -12.83$$

$$k(298 \text{ K}) = 3.0\text{E-}13 \text{ cm}^3/\text{molec.-s.}$$

$$A = 3.2E-12 \text{ cm}^3/\text{molec.-s.}$$

$$E/R = 705 \text{ K}$$

$$\text{-CH}_2\text{F- site: } \log k/n(\text{H}) = -14.79 + G(\text{F}) + G(\text{CH}_3) = -13.02$$

$$k(298 \text{ K}) = 1.9E-13 \text{ cm}^3/\text{molec.-s.}$$

$$A = 3.0E-12 \text{ cm}^3/\text{molec.-s.}$$

$$E/R = 822 \text{ K}$$

The sum of these three site contributions over the temperature range 275 - 425 K corresponds to the following effective Arrhenius parameters:

$$k(\text{total})_{\text{est.}} = 8.9E-12 \exp(-814/T) \text{ cm}^3/\text{molec.-s.}$$

$$k(298 \text{ K})_{\text{est.}} = 5.8E-13 \text{ cm}^3/\text{molec.-s.}$$

*Estimated Rate Constant for CH<sub>3</sub>CHFCH<sub>3</sub> :*

$$2 \text{-CH}_3 \text{ sites: } \log k/n(\text{H}) = -14.79 + G(\text{CH}_2\text{F}) = -14.08$$

(The CH<sub>2</sub>F group has been used as a surrogate for the CH<sub>3</sub>CHF group.)

$$k = 5.0E-14 \text{ cm}^3/\text{molec.-s.}$$

$$A = 6.0E-12 \text{ cm}^3/\text{molec.-s.}$$

$$E/R = 1427 \text{ K}$$

**-CHF- site:**

In this case there are 3 groups to be accounted for; 2 CH<sub>3</sub> groups and one F. Also, this is a case where it makes a difference which group is considered to be the third group, because of the very different G-values between methyl and the F-atom. We therefore assume the three possible cases (two identical) and take the average,

$$\text{(First case) } \log k/n(\text{H}) = -14.79 + G(\text{F}) + G(\text{CH}_3) + 0.27 G(\text{CH}_3) = -12.68$$

$$k = 2.1E-13 \text{ cm}^3/\text{molec.-s}$$

(To be used twice).

$$\text{(Second case) } \log k/n(\text{H}) = -14.79 + 2 \times G(\text{CH}_3) + 0.27 G(\text{F}) = -12.15$$

$$k = 7.1E-13 \text{ cm}^3/\text{molec.-s.}$$

$$k(\text{average}) = 3.8\text{E-}13 \text{ cm}^3/\text{molec.-s.}$$

$$A = 1.9\text{E-}12 \text{ cm}^3/\text{molec.-s}$$

$$E/R = 480 \text{ K}$$

The sum of these two site contributions over the temperature range 275 - 425 K corresponds to the following effective Arrhenius parameters:

$$k(\text{total})_{\text{est.}} = 3.6\text{E-}12 \exp(-633/T) \text{ cm}^3/\text{molec.-s.}$$

$$k(298 \text{ K})_{\text{est.}} = 4.3\text{E-}13 \text{ cm}^3/\text{molec.-s.}$$

The estimation is in good agreement (within 35% at 298 K) with the experimental values from Equations 4 and 5.

The prediction for n-propyl fluoride is probably more reliable than that for isopropyl fluoride, since there is no third group correction in the former case. The predicted 298 K rate constants for the two compounds are similar, but the Arrhenius parameters are quite different. For isopropyl fluoride, about 90% of the reaction occurs at the CHF site, so that the overall reaction has a lower pre-exponential factor, but also a lower activation energy. The n-propyl fluoride has several reactive sites which contribute significantly to the total reaction and therefore lead to a higher pre-exponential factor, but all have relatively high activation energies.

**Acknowledgment.** We have benefited greatly from frequent discussions with members of the JPL Kinetics and Photochemistry group. The DuPont Company generously provided the 281ea sample used in this work. One of us, Edmond Wilson, is very appreciative of being appointed a NASA/ASEE Summer Faculty Fellow at the Jet Propulsion Laboratory, California Institute of Technology for the summers of 1997 and 1998 and to Dr. William DeMore for the opportunity to work with him in his laboratory. This research was carried out by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

## Tables

Table 1. Rate Constant Ratios,  $k_{281ea}/k_{ref}$  as a Function of Temperature Using Infrared Spectroscopy

T(K)	281ea vs Ethane	T(K)	281ea vs Propane	T(K)	281ea vs Ethyl Chloride
288	$2.257 \pm 0.2$	293	$0.605 \pm 0.04$	293	$1.715 \pm 0.04$
289	$2.411 \pm 0.2$	298	$0.556 \pm 0.07$	303	$1.795 \pm 0.04$
298	$2.671 \pm 0.3$	298	$0.598 \pm 0.06$	311	$1.797 \pm 0.05$
298	$2.601 \pm 0.2$	323	$0.510 \pm 0.04$	321	$1.567 \pm 0.06$
298	$2.496 \pm 0.2$	352	$0.523 \pm 0.04$	333	$1.773 \pm 0.02$
308	$2.129 \pm 0.3$	368	$0.513 \pm 0.04$	342	$1.722 \pm 0.02$
315	$2.349 \pm 0.3$	394	$0.434 \pm 0.03$	363	$1.593 \pm 0.03$
328	$1.901 \pm 0.2$			378	$1.590 \pm 0.04$
339	$1.623 \pm 0.1$			386	$1.520 \pm 0.06$
349	$1.460 \pm 0.1$				
358	$1.578 \pm 0.1$				
369	$1.499 \pm 0.1$				
388	$1.304 \pm 0.1$				
394	$1.396 \pm 0.1$				

Table 2: Rate Constant Ratios,  $k_{281ea}/k_{ref}$ , as a Function of Temperature Using Gas Chromatography

T(K)	281ea vs Ethane	T(K)	281ea vs Propane
272	$2.724 \pm 0.08$	270	$0.541 \pm 0.002$
282	$2.529 \pm 0.03$	277	$0.555 \pm 0.003$
298	$2.131 \pm 0.06$	284	$0.523 \pm 0.000$
318	$1.912 \pm 0.03$	292	$0.507 \pm 0.01$
348	$1.601 \pm 0.10$	298	$0.517 \pm 0.01$
361	$1.540 \pm 0.01$	326	$0.496 \pm 0.01$
368	$1.535 \pm 0.04$	352	$0.467 \pm 0.01$
393	$1.358 \pm 0.03$	393	$0.453 \pm 0.01$
413	$1.174 \pm 0.01$	424	$0.432 \pm 0.003$



Table 3. Rate Constants of the Reference Compounds Used in This Work.

Reference Compound	Arrhenius Rate Constant <sup>(a)</sup>	k(298 K) <sup>(a)</sup>	Source
CH <sub>3</sub> CH <sub>3</sub>	1.00E-12 exp(1094/T)	2.54E-13	Atkinson <sup>1</sup>
CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub>	1.29E-11 exp(-730/T)	1.11E-12	DeMore and Bayes <sup>3</sup>
C <sub>2</sub> H <sub>5</sub> Cl	5.14E-12 exp(-855/T)	2.92E-13	Herndon et al. <sup>4</sup>

(a) Units are cm<sup>3</sup>/molec-s

Table 4. Current G-values Used in the Estimation Method.

GROUPS	G - value
F	0.52
2F	0.34
CF3	-0.45
CH2F	0.71
CHF2	-0.07
CH3	1.25
Cl	0.76
2Cl	1.54
CH2Cl	0.76
CH2Br	0.76
CHCl2	0.60
CHFCl	0.05
CF2Cl	-0.19
F,Cl	0.90
CCl3	0.29
CFCl2	0.08
Br	0.79
2Br	1.59
BrCl	1.51
F,Br	1.2
CN	0.52
3rd group multiplier	0.27

Table 5. Pre-exponential Factors for the Reaction of OH with a Variety of Substrates.

<b>Compound</b>	<b>A-Factor</b> cm <sup>3</sup> /molec.-s.
CH <sub>3</sub> CFCl <sub>2</sub> (141b)	1.65E-12
CF <sub>3</sub> CHF <sub>2</sub> CF <sub>3</sub> (227ea)	7.07E-13
CHF <sub>2</sub> Br (22B)	1.53E-12
CHF <sub>2</sub> Cl (22)	9.73E-13
CH <sub>2</sub> F <sub>2</sub> (32)	2.73E-12
CF <sub>3</sub> CH <sub>3</sub> (143a)	1.90E-12
CF <sub>3</sub> CH <sub>2</sub> F (134a)	1.74E-12
CF <sub>3</sub> CH <sub>2</sub> CF <sub>3</sub> (236fa)	1.46E-12
CF <sub>2</sub> HCF <sub>2</sub> H (134)	2.05E-12
CHBr <sub>3</sub>	1.36E-12
CF <sub>2</sub> ClCCl <sub>2</sub> H(122)	8.50E-13
CFCI <sub>2</sub> CHFCl (122a)	1.08E-12
CH <sub>3</sub> Cl	7.46E-12
CH <sub>3</sub> F	4.20E-12
CH <sub>3</sub> Br	6.90E-12
CHCl <sub>3</sub>	1.80E-12
CF <sub>3</sub> OCH <sub>3</sub> (143aE)	2.98E-12
CHCl <sub>2</sub> CF <sub>3</sub> (123)	7.75E-13
CF <sub>3</sub> CF <sub>2</sub> CF <sub>2</sub> OCHF <sub>2</sub> CF <sub>3</sub>	7.44E-13
CF <sub>3</sub> OCHF <sub>2</sub> CF <sub>3</sub> (227eaE)	7.09E-13
CH <sub>3</sub> OCH <sub>3</sub>	1.49E-11
c-C <sub>6</sub> H <sub>12</sub>	3.71E-11
c-C <sub>5</sub> H <sub>10</sub>	2.57E-11
c-C <sub>3</sub> H <sub>6</sub>	5.15E-12
c-C <sub>4</sub> H <sub>8</sub>	1.62E-11

## References

- (1) Atkinson, R. *J. Phys. Chem. Ref. Data* **1997**, *26*, 215-290.
- (2) DeMore, W. B. *J. Phys. Chem.* **1996**, *100*, 5813-5820.
- (3) DeMore, W. B.; Bayes, K. D. *J. Phys. Chem.* **1998**, submitted.
- (4) Herndon, S.; Gierczak, T.; Talukdar, R. K.; Ravishankara, A. R. **1998**, In preparation.
- (5) Hsu, K. J.; DeMore, W. B. *J. Phys. Chem.* **1995**, *99*, 1235-1244.
- (6) Hsu, K. J.; DeMore, W. B. *J. Phys. Chem.* **1995**, *99*, 11141-11930.

## Figure Captions

Figure 1. Rate constant data for HFC-281ea, based on the ratios as measured by GC vs ethane and propane.

Figure 2. Comparison of the IR results with the Arrhenius line representing the GC data.

Figure 3. A plot of A-factors for several reactions vs the log of the rate constant per H-atom.

Figure 1




Figure 2




Figure 3

