

S. D. Lord(1), S. J. Unger(2), J. Brauher(1), P. E. Clegg(2), P. Cox(3),  
 J. Fischer(4), M. Greenhouse(5), M. L. Luhman(4), M. A. Malkan(6),  
 S. Satyapal(5), H. A. Smith(7), L. Spinoglio(8), G. J. Stacey(9), M. Wolfire(10)

- 1 California Institute of Technology, Pasadena, USA
- 2 Queen Mary and Westfield College, London, UK
- 3 Institut d'Astrophysique Spatiale, France
- 4 Naval Research Laboratory, Washington, USA
- 5 NASA Goddard, Greenbelt, USA
- 6 University of California, Los Angeles, USA
- 7 Smithsonian Astrophysical Observatory, Cambridge, USA
- 8 Istituto di Fisica dello Spazio Interplanetario-CNR, Frascati, Italy
- 9 Cornell University, Ithaca, NY, USA
- 10 University of Maryland, College Park, USA

## Abstract

We have obtained low resolution ( $R=200$ ) fully sampled far-infrared (FIR) 42-197  $\mu\text{m}$  spectra of the southern AGN/starburst SB(s)c galaxy NGC 4945 with the Long Wavelength Spectrometer (LWS) on the Infrared Space Observatory (ISO). The galaxy was observed at a central, NE, and SW position, each separated by 165" with the instrument's  $\sim 75$ " aperture. The forbidden fine-structure transitions of [CII] 158  $\mu\text{m}$  and [OI] 63  $\mu\text{m}$  lines were detected at each position. Detections of the [OI] 145  $\mu\text{m}$  and [OIII] 88  $\mu\text{m}$  lines were also obtained at some positions. Using these lines and the total FIR continuum flux, we have modeled the properties of the warm atomic gas with photodissociation region (PDR) models (Tielens and Hollenbach, 1985; Wolfire et al. 1990; Kaufman et al. 1998). The low-J rotational transitions of OH at 53  $\mu\text{m}$  and 119  $\mu\text{m}$  are seen in absorption at the nuclear position. The total OH column density and the role of these lines in the pumping of the OH radio maser are considered for these two transitions. The NE and SW positions lie on the massive molecular ring. The ring and the nuclear conditions are analyzed by comparing the CII/OI, CII/CO, and (CII+OI)/FIR ratios.

NGC4945 is thought to have a combination AGN and starburst nucleus, and also displays dramatic FIR and CO hot--spots at  $\sim 180''$  from the nucleus. Our ISO observations show strong [OI] 63  $\mu\text{m}$  and [CII] 158  $\mu\text{m}$  fine structure emission lines at all three positions. These may be result of the heating of interstellar clouds by massive stars. The nuclear position shows a very low CII line/total FIR-continuum ratio (less than 0.1 %). More typical line/continuum ratios ( $\sim 0.5$ ) are seen in the hot--spots. All three positions show [OI]/[CII] ratios near unity.

From our observations, gas heating in the nuclear position from a central AGN can not be ruled-out. (Shorter wavelength, higher excitation FIR transitions observed by ISO will be used to help define the role of an AGN here). If the dominant heating nuclear source is from massive star formation, then strong interstellar UV radiation fields,  $3 \times 10^4$  times the typical Milky Way field, are required to produce the line/continuum ratios seen. These are high field strengths even for starburst galaxies. The NE and SW hot-spot positions show rather typical starburst UV fields,  $3 \times 10^3$  times the Milky way value, and somewhat higher atomic gas densities ( $n \sim 10^4$ ).

The clouds in these three regions are small,  $R \sim 1-10\text{pc}$ , have warm neutral gas surfaces,  $T \sim 200\text{K}$ , and are concentrated with high volume filling factors in the nucleus (0.5) and smaller filling factors in the hot--spots:  $\sim 10^{-4}$ . In the nucleus, either there are several clouds along a line of sight, or the gas is arranged in sheets or filaments.

In the nucleus, five transitions of OH are seen (mostly absorption lines) and these provide a consistent excitation and de-excitation picture for the low--level energy. Also in the nucleus we observe two transitions of ortho-water in absorption: at 101  $\mu\text{m}$  and 179  $\mu\text{m}$ .