

# Observation of Global Ocean-Atmosphere Exchanges from Spaceborne Sensors

W. Timothy Liu

Jet Propulsion Laboratory 300-323

California Institute of Technology

Pasadena, CA 91109, U.S.A.

(Tel: 1-818-354-2394; Fax: 1-818-393-6720; email:liu@pacific.jpl.nasa.gov)

## INTRODUCTION

The ocean and the atmosphere are coupled by the fluxes of momentum, heat, and water, but in situ measurements of these fluxes are sparse and uneven. Global fields of these fluxes can be derived from satellite data or obtained from operational numerical weather prediction (NWP) models. The NWP models provide quality control and dynamic interpolation of sparse and uneven data. However, dynamic interpolation is only as good as the model physics and parameterization scheme, and over most of the ocean, there may not be sufficient data for any useful interpolation. operational NWP models change frequently; the results are inconsistent of climatic studies. Reanalysis to produce long period of consistent data have only been started recently. Adequate observations at temporal and spatial scales significant to global change research can only be achieved from the vantage point of space. Spaceborne sensors provide repeated global observations of some parameters from which these fluxes can be derived. The status and shortfalls of spacebased estimation of the fluxes will be summarized and assessed in this report. Satellite data can be assimilated into numerical model to simulate surface fluxes, but the progress in developing assimilation technique of unconventional satellite data has been slow.

## MOMENTUM FLUX

The flux of momentum and kinetic energy are resulted from wind shear. There are three sets of spacebased global wind field available now and in the near future. The Special Sensor Microwave Imager (SSM/I) on the operational spacecraft of the Defense Meteorological Space Program has provided continuous wind speed measurements over global ocean since July 1987 [1]. It has a wide scan and, therefore, good coverage, but lacks directional information. The wind speed from SSM/I has been combined with other available wind data through a variation method to produce wind vector fields at 6 hourly interval and 2° by 2.5° resolution. These wind fields were found to generate more realistic anomalous ocean cooling in an ocean general circulation model [2]. SSM/I is an operational sensor and data availability will be continued well into the next century. The production of wind vector, however, is a research effort.

There are scatterometers on board European Remote Sensing Satellites (ERS) which were specifically designed to measure surface wind vector [3]. ERS-1 was launched in 1991 with a C-band scatterometer and it was replaced by a duplicate on ERS-2 in 1996. Unfortunately the ERS scatterometers scan a narrow 475 km swath only on one side of the spacecraft and leaves large data gaps between passes; the coverage is less than half of SSM/I. ERS winds from January 1992 have been available but the distribution is restricted. The next European scatterometer will be on the Meteorology Operational Platform (METOP) to be launched in 2002.

The NASA scatterometer (NSCAT) was launched in August 1996 on the Japanese spacecraft ADEOS-1. It will scan two 600 km swaths on both sides and will have more than twice the coverage of ERS scatterometers. It will cover over 87% of the ocean in one days, and 97% in two days, under both clear and cloudy conditions. NSCAT measures at Ku-band which has been studied more vigorously than C-band. NSCAT will be followed by an improved version, called SeaWinds, on ADEOS-2 to be launched in 1999. A third scatterometer is being proposed for ADEOS-3 to be launched in 2002. The third version will incorporate a polarimetric radiometer as a proof-of-concept study of measuring wind vector with a passive sensor. Spacebased observations of wind vector will be continued on the National Polar-orbiting Operational Satellite System (NPOESS) in the next century. Summaries of scatterometry and examples of applications are given in [4].

## HEAT FLUX

Air-sea heat flux can be divided into four components; sensible heat resulted from thermal gradient, latent heat (LF) carried by evaporation, shortwave radiation (SR) from the sun and longwave radiation from the atmosphere and the ocean. Shortwave radiation and latent heat flux are the larger variable components over most of the tropical and temperate oceans.

Global SR has been computed from the Earth Radiation Budget Experiment (ERBE) data from 1985 to 1989 [5]. The variability of SR is largely controlled by the variability of

clouds and most of the computations of SR make use of the high resolution and high sampling of data from **geostationary** satellites. The International Satellite Cloud Climatology Project (ISCCP) has provided calibrated and standardized cloud data from 4 **geostationary** satellites operated by the U.S., the Japanese, and the European space agencies. A number of methods [6, 7, 8] have been used to compute SR at daily **and**  $2^\circ$  by  $2.5^\circ$  resolution, including the operational effort by the Surface Radiation Budget Program at **Langley** Research Center. The availability of ISCCP data is expected well into the future. The launching of sensors for the Earth Observation Program, including CERES (Clouds and Earth Radiant Energy System) will advance the derivation of SR.

The computation of LF requires sea surface temperature (SST), wind speed ( $u$ ), and humidity at a level within the surface layer ( $q$ ). Over ocean,  $u$  and SST have been directly retrieved from satellite data, but not  $q$ . A method of estimating  $q$  and LF from the ocean, using satellite data was proposed [9] which is based on an empirical relation between the integrated water vapor (measured by spaceborne microwave radiometer) and  $q$  (required to compute evaporation) in the **monthl**y time scale [10]. The physical rationale is that the vertical distribution of water vapor through the whole depth of atmosphere is coherent for period longer than a week [11]. The relation does not work well at synoptic and shorter time scales and also fails in some regions during summer [12]. The relation have also been scrutinized **in a number of studies** [13, 14, 15]. Methods which include other geophysical parameters or EOF functions as estimators have also been proposed [16, 17, 18, 19]. Future advanced microwave humidity sounders may improve the accuracy and **temporal** resolution of evaporation estimation.

It is much more difficult to estimate sensible heat flux and **longwave** radiation from satellite data. The **magnitude** and variability of sensible heat are relatively small over much of the ocean and near surface air temperature is difficult to retrieve with sufficient accuracy. It can be derived from LF if the relative humidity or the Bowen Ratio is known, but the accuracy is uncertain [20]. Longwave radiation is strongly **affected** by atmospheric properties below cloud base which are hidden from spaceborne sensors. Methods that combined cloud information with atmospheric soundings and numerical models have been attempted [21, 22]. **Improved** atmospheric temperature sounders such as Atmospheric Infrared Sounder (AIRS), scheduled to be launched early next decade may advance the estimation of these fluxes.

## WATER FLUX

**Hydrologic** forcing is the **difference** between precipitation and evaporation. The estimation of evaporation is the same

as the estimation of latent heat flux discussed above. Attempts have also been made to estimate precipitation from **spaceborne** sensors at visible, infrared, and microwave wavelengths [23, 24]. Passive microwave sensors have the advantage of more directly related to rain but are limited by the insufficient sampling of the diurnal cycle by the **polar** orbiters. Data production and validation have been **undertaken** by the Global **Precipitation** Climatology Project (GPCP) and data at  $2.5^\circ$  and monthly resolutions from 1986 to 1994 **are** available. Precipitation for a very long **period** of time has also been **derived** from the operational Microwave Sounding Unit [25]. Improvement in rain measurement is **expected** with the launching the Tropical Rain Measuring Mission (TRMM) in 1997.

## ASSESSMENT

At present, we have the ability of monitoring momentum flux over global ocean using **spacebased** data. We **could** discern the annual variations over a large part of the global ocean and El Niño anomalies over the tropical ocean of the two major components of heat flux. However, we **do** not have the absolute accuracy of the net heat flux to compute the average meridional ocean heat transport from **spacebased** heat flux estimates. The retrieving method of **q** **does** not work well at high frequencies and over some region. The validation of **spacebased** estimation of precipitation has been hampered by the lack accurate standard. Surface **wind** measurement is sustainable if **the** series of planned **scatterometers** are approved. While present methods of estimating heat and water fluxes rely on operational sensors which are sustainable, the estimation of the fluxes from the satellite **data** are largely research efforts that need to be supported, before more stable and operational methods emerge.

The studies of air-sea fluxes involve the interaction of the atmospheric and ocean and are **intrinsically** interdisciplinary. Spacebased estimation of these fluxes required synergistic combination of multiple sensors from different flight projects. To pursue such studies, traditional disciplinary separations (in the research institutes or in the funding agencies) have to be overcome and individual flight project prepossession also has to **be challenged**. Coordination in sensor deployment and data processing is **needed** to have coincident or continuous data. The limited life-span of a single sensor requires continuous sensor deployment to study **interannual** to **decadal** changes. Because spaceborne sensors are expensive and require long lead time to develop, continuous vigilance from the scientific community to advocate and coordinate is necessary to shepherd these space programs to fruition.

## ACKNOWLEDGMENTS

This study was performed at the Jet Propulsion Laboratory, under contract with the National Aeronautics and Space Administration (NASA). It was supported by the Earth Observing System Interdisciplinary Science Program and the Physical Oceanography Program of NASA's Mission to Planet Earth.

#### REFERENCES

- [1] Wentz, F.J., "Measurement of oceanic wind vectors using satellite microwave radiometer", *IEEE Trans. Geosci. Remote Sens.*, 30, 960-972, 1992.
- [2] Liu, W.T., W. Tang, and R. Atlas, "Responses of the tropical Pacific to wind forcing as observed by spaceborne sensors and simulated by an ocean general circulation model", *J Geophys. Res.*, 101, 16345-16359, 1996.
- [3] Attema, E.P.W., "The active microwave Instrument on board the ERS-1 satellite", *Proc. IEEE*, 79, 791-799, 1991.
- [4] Liu, W.T. and W. Tang, "Spaceborne scatterometer in studies of atmospheric and oceanic phenomena from synoptic to interannual time scales". *Space Remote Sensing of Subtropical Ocean*. C.T. Liu (cd.), Elsevier Press, Amsterdam, in press.
- [5] Li, Z, and H.G. Leighton, "Global climatologies of the solar radiation budgets at the surface and in the atmosphere from 5 years of ERBE data.", *J. Geophys. Res.*, 98, 4919-4930, 1993.
- [6] Gautier, C., G. Disk, and S. Masse, "A simple physical model to estimate incident solar radiation at the surface from GOES satellite data" *J. Appl. Meteor.*, 19, 1821-1844, 1980.
- [7] Bishop, J.K.B., and W.B. Rossow, "Spatial and temporal variability of global surface solar irradiance" *J. Geophys. Res.*, 96, 16839-16858, 1991.
- [8] Pinker, R.T., and I. Laszlo, "modeling surface solar irradiance of satellite applications on a global scale", *J. Appl. Meteor.*, 31, 194-211, 1992.
- [9] Liu, W.T., and P.P. Niiler, "Determination of monthly mean humidity in the atmospheric surface layer over oceans from satellite data", *J. Phys. Oceanogr.*, 14, 1451-1457, 1984.
- [10] Liu, W.T., "Statistical relation between monthly precipitable water and surface-level humidity over global oceans. *Mon. Wea. Rev.*, 114, 1591-1602, 1986.
- [11] Liu, W.T., W. Tang, and P.P. Niiler, "Humidity profiles over ocean", *J. Clim.*, 4, 1023-1034, 1991.
- [12] Liu, W.T., W. Tang, and F. J. Wentz, "Precipitable water and surface humidity over global oceans from SSM/I and ECMWF", *J. Geophys. Res.*, 97, 2251-2264, 1992.
- [13] Hsu, S.A., and B.W. Blanchard, "The relationship between total precipitable water and surface-level humidity over the sea surface: a further evaluation", *J. Geophys. Res.*, 94, 14539-14545, 1989.
- [14] Esbensen, S.K., D.B. Chelton, D. Vickers, and J. Sun, "An analysis of errors in Special Sensor Microwave Imager evaporation estimates over the global oceans" *J. Geophys. Res.*, 98, 7081-7101, 1993.
- [15] Jourdan, D. and C. Gautier, "Comparison between global latent heat flux computed from multi-sensor (SSM/I and AVHRR) and from in-situ data", *J. Atmos. and Oceanic Tech.*, 12, 46-72, 1994.
- [16] Wagner, D., E. Ruprecht, and C. Simmer, "A combination of microwave observations from satellite and an EOF analysis to retrieve vertical humidity profiles over the ocean", *J. Appl. Meteor.*, 29, 1142-1157, 1990.
- [17] Cresswell, S., E. Ruprecht, and C. Simmer, "Latent heat flux over the North Atlantic Ocean-A case study. *J. Appl. Meteor.*, 30, 1627-1635, 1991.
- [18] Miller, D.K., and K.B. Katsaros, "Satellite-derived surface latent heat fluxes in a rapidly intensifying marine cyclone" *Mon. Wea. Rev.*, 120, 1093-1107, 1991.
- [19] Chou, S.H., R.M. Atlas, and J. Ardizzone, "Estimates of surface humidity and latent heat fluxes over oceans from SSM/I data", *Mon. Wea. Rev.*, 123, 2405-2435, 1995.
- [20] Liu, W.T., and P.P. Niiler, 1990: The sensitivity of latent heat flux to the air humidity approximations used in ocean circulation models. *J. Geophys. Res.*, 95, 9745-9753, 1990.
- [21] Frouin, R., C. Gautier, and J. Morcrette, 1988: Downward longwave irradiance at the ocean surface from satellite data: methodology and in situ validation. *J. Geophys. Res.*, 93, 597-619.
- [22] Gupta, S.K., W. Darnell, and A.C. Wilber, 1992: A parametrization for longwave surface radiation from satellite data: recent improvements. *J. Appl. Meteor.*,
- [23] Arkin, P.A., P.E. Ardanuy, "Estimating climatic-scale precipitation from space: a review", *J. Climate*, 2, 1229-1238, 1989.
- [24] Wilheit, T.T., A.T.C. Chang, and L.S. Chiu, "Retrieval of monthly rainfall indices from microwave radiometric measurements using probability distribution functions", *J. Atmos. Oceanic Tech.*, 8, 118-136, 1991.
- [25] Spencer, R.W., "Global oceanic precipitation from the MSU during 1979-92 and comparisons to other climatologies", *J. Clim.*, 6, 1301-1326, 1993.