

## THE HOT BANDS OF METHANE BETWEEN 5 AND 10 MICRONS

O. Ouardi, J. C. Hilico, M. Loete, *Laboratoire de Physique de l'Université de Bourgogne*, and L. R. Brown, Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, California 91109.

The experimental line intensities of 1727 transitions between 1000 and 2000  $\text{cm}^{-1}$  belonging to the pentad-dyad system of methane are fitted to first and second order using the effective dipole moment expansion in the polyad scheme. The nine observed bands are  $\nu_3-\nu_2$ ,  $\nu_3-\nu_4$ ,  $\nu_1-\nu_2$ ,  $\nu_1-\nu_4$ ,  $2\nu_4-\nu_4$ ,  $2\nu_2-\nu_4$ ,  $2\nu_2-\nu_2$ ,  $\nu_2+\nu_4-\nu_2$  and  $\nu_2+\nu_4-\nu_4$ . The intensities are obtained from long-path spectra recorded at  $0.005\text{ cm}^{-1}$  resolution using the McMath Fourier transform spectrometer located at Kitt Peak National Observatory. To achieve a more stable fit for three of the 27 constants required in the second order model, 1206 intensities of the dyad ( $\nu_2$ ,  $\nu_4$ ) are refitted simultaneously with the hot band data. The fits to first and second order lead to rms values, respectively, of 21.5% and 5.0% for the 1727 hot band intensities and 6.5% and 3.0% for the 1206 dyad lines. The band strengths of all 10 pentad-dyad hot bands range from 0.93 for  $2\nu_4-\nu_4$  to  $7.7 \times 10^{-5}$  for the unobserved  $2\nu_4-\nu_2$  in units of  $\text{cm}^{-2}/\text{atm}$  at 296 K.

Part of the research reported in this paper was performed at the Jet propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

---

Time required: 15 min

Session in which paper is recommended for presentation: 4