

# ELECTRON IMPACT IONIZATION AND DISSOCIATIVE IONIZATION OF C<sub>2</sub>H<sub>2</sub>

Shuanghai Zheng\* and S. K. Srivastava

Jet Propulsion Laboratory, California Institute of technology, 4800 Oak Grove Dr., Pasadena, California, 91109, USA

By utilizing a crossed electron beam collision geometry, a combination of time-of-flight (TOF) and quadrupole mass spectrometers, and the relative flow technique<sup>1</sup> normalized values of cross sections and appearance energies (AP) were obtained for the formation of singly and multiply ionized species resulting from the ionization and dissociation of C<sub>2</sub>H<sub>2</sub>. Details on the apparatus and technique have been published previously<sup>2,3</sup>.

Figure 1 shows spectra of ions obtained at 100 eV electron impact energy. In this figure those species are shown for which "ionization efficiency curves" were acquired in the energy range from threshold to 800 eV. The shapes of these ionization efficiency curves were corrected by comparing to similar curves for the rare gas atoms. From these curves the AP for each species was derived. Finally, the ionization efficiency curves were employed to generate normalized values of cross sections by utilizing the relative flow technique and known cross sections for H<sub>2</sub>, Ne, and N<sub>2</sub> at 100 and 250 eV electron impact energies. Previous measurements on this molecule are by Tate et al.<sup>4</sup> and Hangmann<sup>5</sup>. For the case of the formation of C<sub>2</sub>H<sub>2</sub><sup>+</sup> they are shown in fig. 2. For C<sub>2</sub>H<sub>2</sub><sup>+</sup>, C<sub>2</sub>H<sup>+</sup>, and C<sub>2</sub><sup>+</sup> there is a reasonable agreement between the various results. However, the differences are large in the cross section values for the formation of C<sub>2</sub>H<sub>2</sub><sup>+</sup>, C<sub>2</sub><sup>++</sup>, and H<sup>+</sup>.

AP's for various ions have also been measured previously by Tate et al.<sup>4</sup> which agree very well with the present values. These AP's were helpful in identifying the various pathways for the dissociation of C<sub>2</sub>H<sub>2</sub> into ionic species.

\* NRC/NASA Research Associate

## Reference

1. S. K. Srivastava, A. Chutjian, and S. Trajmar, J. Chem. Phys. **63**, 2659(1975).
  2. E. Krishnakumar and S. K. Srivastava, J. Phys. B: At. Mol. Opt. Phys. **21**, 105 S(1988).
  3. E. Krishnakumar and S. K. Srivastava, J. Phys. B: At. Mol. Opt. Phys. **23**, 1893(1991).
- It is well
4. J. T. Tate, P. T. Smith and A. L. Vaughan, Phys. Rev. **48**, 525(1935).
  5. S. A. Gaudin and R. Hagemann, J. Chim. Phys. **64**, 1209(1976).




Fig. 1. Spectra of ions resulting from C<sub>2</sub>H<sub>2</sub> at 100 eV electron impact energy.




Fig. 2. Cross sections for the formation of C<sub>2</sub>H<sub>2</sub><sup>+</sup>.