

***HST* Observations of the Luminous IRAS Source FSC10214+4724:
A Gravitationally Lensed Infrared Quasar'**

Peter R. Eisenhardt², Lee Armus³, David W. Hogg³, B. T. Soifer³, G. Neugebauer³,
Michael W. Werner²

Received _____, accepted _____

¹Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by AURA under NASA contract NAS 5-26.555.

²MS 169-327, Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109

³Division of Physics, Math, & Astronomy, California Institute of Technology, Pasadena, CA 91125

ABSTRACT

With a redshift of 2.3, the IRAS source FSC10214+4724 is apparently one of the most luminous objects known in the Universe. We present an image of FSC10214+4724 at $0.8\mu\text{m}$ obtained with the Hubble Space Telescope WFPC2 Planetary Camera. The source appears as an unresolved ($< 0''.06$) arc $0''.7$ long, with significant substructure along its length. The center of curvature of the arc is located near an elliptical galaxy $1''.1$ S to the north. An unresolved component 100 times fainter than the arc is clearly detected on the opposite side of this galaxy. The most straightforward interpretation is that FSC10214+4724 is gravitationally lensed by the foreground elliptical galaxy, with the faint component a counterimage of the IRAS source. The brightness of the arc in the *HST* image is then magnified by ~ 100 and the intrinsic source diameter is $\sim 0''.01$ (50 pc) at $0.25\mu\text{m}$ rest wavelength. The bolometric luminosity is probably amplified by a smaller factor (~ 30), due to the larger extent expected for the source in the far-infrared. A detailed lensing model is presented which reproduces the observed morphology and relative flux of the arc and counterimage, and correctly predicts the position angle of the lensing galaxy. The model also predicts reasonable values for the velocity dispersion, mass, and mass-to-light ratio of the lensing galaxy for a wide range of galaxy redshifts. A redshift for the lensing galaxy of ~ 0.9 is consistent with the measured surface brightness profile from the image, as well as with the galaxy's spectral energy distribution. The background lensed source has an intrinsic luminosity $\sim 2 \times 10^{13} L_{\odot}$, and remains a highly luminous quasar with an extremely large ratio of infrared to optical/ultraviolet luminosity.

Subject headings: infrared: galaxies, cosmology: gravitational lensing,
galaxies: individual

1. Introduction

Ever since its identification with a redshift 2.286 optical emission line source by Rowan-Robinson et al. (1991), leading to an inferred bolometric luminosity $\sim 5 \times 10^{14} L_{\odot}$, the IRAS source FSC1 0214+4724 has been the subject of enormous attention. Detections of CO (Brown & VandenBout 1991; Solomon, Downes & Radford 1992; Tsuboi & Nakai 1992) and submillimeter continuum emission (Clements et al. 1992, Downes et al. 1992) from the source confirmed the presence of huge quantities of gas and dust. With a vastly larger lookback time and luminosity than any other known IRAS source, FSC10214+4724 appeared to be either an extremely luminous dust embedded quasar, or a representative of a new class of astronomical object, e.g. a primeval galaxy.

However, while the redshift of the IRAS source is secure, its intrinsic luminosity is less certain. The fact that FSC1 0214-E4724 lies at the flux limit of the IRAS survey, combined with the presence of several red companion objects within a few arcseconds, led Elston et al. (1994) to suggest that the IRAS source might be gravitationally lensed by a foreground group of galaxies. Intriguingly, Matthews et al. (1994) found arcs emerging from the source in a deconvolved K band image with $0''.6$ seeing taken with the Keck telescope. Matthews et al. considered the lensing hypothesis, but concluded it was unlikely because the image morphology was not achromatic. Broadhurst & Lehar (1995) modelled the source as gravitationally lensed, finding support for their model from a reanalysis of the Matthews et al. data. Graham & Liu (1995) also argue for lensing, based on deconvolution of a more recent (March 1995) Keck K band image with $0''.4$ seeing. Trentham (1995) argues on statistical grounds that the probability of a unification due to lensing is likely to be less than a factor of ten.

We present an image of FSC1 0214+4724 taken in December 1994 at 8000 \AA with the *HST* WFPC2 Planetary Camera with $0''.1$ resolution. This image provides dramatic support for the lensing hypothesis, implying a magnification in the *HST* data of ~ 100 . We

use the image to derive a detailed model for the intrinsic properties of the lensed source and the lensing galaxy.

For reference, at the FSC 10214+4724 redshift $z = 2.286$, one $0''.0455$ Planetary Camera pixel subtends $300(180)h^{-1}\text{pc}$ for $q_0 = 0(0.5)$, while these values are $239(191)h^{-1}\text{pc}$ for $z = 0.9$, where $h \equiv H_0 / 100 \text{ km sec}^{-1} \text{ Mpc}^{-1}$. Where not otherwise specified, we assume $H_0 = 50 \text{ km sec}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0.5$.

2. Observations and Reduction

Three frames, each 2200 seconds long, were obtained on 11 consecutive orbits with the WFPC2 F814W filter on the 10th and 11th of December 1994 (UT). FSC 10214+4724 was positioned near the center of the Planetary Camera, and each exposure was displaced from the other two by an integer number (5 or 10) of PC pixels in both axes. The Wide Field Camera data are not considered here.

After standard processing provided by STScI, the multiple frames were used to filter out cosmic rays and hot pixels. Although these defects are quite prominent and affect roughly 4% of the pixels in each frame, the main characteristics of the combined image discussed in section 3.1. are discernible in each frame even without this filtering.

Cross-correlations were performed on pairs of frames to confirm that the actual displacements between frames, as measured in pixels, were integers to within 0.2 pixels. The frames were then trimmed by the appropriate number of rows and columns to coregister them, and the STSDAS task CRRED was used to average them together, iteratively excluding pixels which deviated from the previous iteration's average value by more than three sigma. The minimum value at each pixel location was used for the initial estimate of the average, and sigma was the value expected from Poisson statistics and the gain and read

noise. To remove multiple pixel cosmic ray events, a stricter limit of 1.5 sigma was applied to the four pixels adjacent to any pixel which exceeded the three sigma criterion. Finally a median filtering routine was applied to identify and interpolate over a few dozen isolated pixels which deviated sharply from their neighbors in the average image, presumably because they were corrupted in all three frames. None of these latter pixels fall within objects in the field, and only a handful of the pixels in the components discussed below are based on data from less than two frames.

3. Results

The combined image of the full Planetary Camera field is shown in Figure 1 (a), while figures 1 (b) and (c) show the FSC10214+4724 region in progressively greater detail.

A synthetic point spread function (PSF) derived from the "Tiny Tim" *HST* image modelling software package was used to deconvolve the average image, because a good empirical point spread function was not available (see section 3.2.). The synthetic PSF was calculated for a source with the color of a I{-star in 14'81.4W at the location of FSC10214+4724 in the Planetary Camera field. Figure 1 (d) shows the same region covered in figure 1 (c) after a mild deconvolution of the data (10 iterations of the STSDAS implementation of the Lucy-Richardson algorithm) onto a grid subsampled four times more finely than the original pixels.

3.1. Morphology

At the resolution of the Planetary Camera, an arc-like structure dominates the morphology of the emission line source, in the terminology of Matthews et al. (1994), which is adopted here, the arc-like structure is component 1 (see Fig. 1 (h)). The extent of this

arc is smaller than shown in Matthews et al., and there is a sharply defined ridge of high surface brightness emission which is $0''.7$ long and essentially unresolved in the transverse direction. Lower surface brightness emission can be seen extending the arc $\sim 0''.4$ to the west, and a similar amount (but at a considerably fainter level) to the east-northeast. There is also a hint of still fainter emission extending a few tenths of an arcsecond due east (*not* along a circular arc) from the eastern tip of the bright ridge. Within the bright ridge are at least two peaks separated by $0''.24$, with the brighter peak towards the east. The center of curvature of the arc was fitted and found to be $\sim 0''.12$ west-northwest, of the center of component 2 (which is $1''.18$ from the arc). Component 2 has a smooth light distribution which is resolved and slightly elongated (see sections 3.2. and 4.3.). Directly opposite component 2 from the arc is a faint but clearly visible source (component 5 in figure 1 (b)), $0''.43$ from the center of component 2. Component 3 is resolved and has a feature which is suggestive of a tidal arm leading back towards component 2. Component 4 appears to be a highly inclined galaxy.

3.2. Brightness Profiles

In an attempt to quantify the radial extent of the arc, pixels from the sector subtended by the brightest $0''.5$ of the arc, at component 2 were sorted in order of radius from component 2. To reduce the effect of the tangential substructure along the arc, a running average of the flux from 5 pixels in this radially sorted list was calculated. Figure 2 plots this running average flux as a function of the average radius of those pixels less the $1''.18$ distance of component 1 from component 2. For comparison, the (unaveraged) radial profiles are plotted for stars A and 11 (see figure 1 (a)), components 2 and 5, and the synthetic PSF which was used in the deconvolution shown in figure 1 (d).

While the wings of the synthetic PSF fall inside those of the arc, the *empirical* PSFs of

stars A (outside its saturated core) and 11 match the arc cross section reasonably well. It therefore appears likely that the synthetic PSF under estimates the FWHM of the true PSF. Based on the synthetic PSF we estimate an upper limit of $0''.06$ (500 pc) for the intrinsic FWHM of the arc in the radial direction. Note that the effects of the running average, of any error in using component 2 as the center of the arc, and of the smaller size of the synthetic PSF all work in the direction of leading us to overestimate this dimension.

The deconvolved image shown in Figure 1 (c) also yields a $0''.06$ FWHM for the arc, but this holds true for star H after deconvolution as well. Because the individual frames are separated by integer numbers of PC pixels, there is little leverage on finer scale structure. Deconvolution does emphasize the high surface brightness of the arc however, increasing it by a factor of three.

In short, we see no evidence that component 1 is resolved in the radial direction. (In section 4.1. we will argue that the intrinsic FWHM of the arc is $\sim 0''.01$). Component 5 also appears unresolved, although its profile suffers from much lower signal to noise.

Component 2, however, is clearly resolved in figure 2. To extend the measurement of component 2's surface brightness profile to larger radii the image was rotated 180° about the center of component 2, and pixels at the locations of other objects in the original image were replaced with pixels from the rotated image. This assumes elliptical symmetry for component 2 in the replaced regions, which cover a maximum of 25% (at $r = 1''.3$) of the area at any radius, and 7% of the total area. Figure 3 shows the resulting radial surface brightness profile for component 2. A deVaucouleurs' profile with an effective radius $r_e \approx 1''.3$ (10 kpc) provides a much better fit to component 2 than do exponential disk models, suggesting that this object is an early type galaxy. The measured ellipticity of component 2 inside the arc is $\approx 0.16 \pm 0.1$ at a position angle of $\approx 3 \pm 15^\circ$ east of north. Excess surface brightness appears near a radius of $1''.4$ even though the component 1 pixels

(which are near this radius) have been replaced. As a check, the surface brightness profile was measured within sectors centered on component 2 from position angles $73-133^\circ$ and $233-318^\circ$, angles which bypass all obvious emission sources in figure 1(b). The value for r_e in this case was $1''.0$ (a smaller r_e is consistent with these sectors being along the minor axis), and excess light, was again found near $1''.4$ radius. The total excess light at this radius is very roughly equivalent to a 23rd magnitude source.

3.3. Photometry

Photometric measurements obtained from the Planetary Camera image for the components are given in Table 1. One count in the image corresponds to 1.185×10^{-21} erg/cm²/sec/Å or to a magnitude of 30.00 in the F814W band with Vegaset to magnitude 0. From the measured standard deviation per pixel, the sensitivity limit (3σ) is $m_{814} \sim 28.2$ mag for a point source or $\mu_{814} \sim 25.6$ mag arcsec⁻². Positions are relative to component 2, whose position in the 11 ST guide star system is given in Table 1. Polygonal apertures were used to include the faint emission seen extending from components 1 and 3. The flux for component 5 was measured using a $0''.35$ diameter aperture, with the local background measured using the mode of an annulus of width $0''.1$ surrounding this aperture, and corrected for PSF losses using the star 11 curve of growth. This flux was checked by subtracting away the image rotated 180° about component 2, and also by subtracting the elliptical model fit to component 2 discussed in section 3.2.. All three methods consistently gave a value close to 100 for the flux ratio of component 1 to component 5, and we adopt 100 for this important ratio for the remainder of the paper.

4. Discussion

The morphology of the components of FSC10214+4724, a circular arc (component 1) with its radius of curvature centered on another object, (component 2), and another fainter image (component 5) on the opposite side, strongly supports the gravitational lens hypothesis, i.e. that component 2 is a foreground galaxy and components 1 and 5 are images of a single background object. The multiple imaging and the arclike morphology and high inferred luminosity of component 1 result from distortion and magnification by the gravitational potential of the foreground component 2. Components 3 and 4 are other galaxies along the line of sight, possibly related to the galaxy which is component 2, and probably involved in the lensing.

The high resolution of the *HST* image makes the arc morphology and component 5 readily apparent, and allows us to directly measure the ratio of the brightnesses of these components. This morphology and ratio are crucial elements in the development of a lens model for the source. We find additional support for the lens hypothesis from the observed morphology of component 2, in particular, as shown in Appendix A, component 2 has the surface brightness profile and spectral energy distribution expected for a foreground elliptical galaxy, and its position angle is correctly predicted by the lens model. In the following, we adopt the interpretation of FSC10214+4724 as a gravitationally lensed system, and describe the detailed model of this system and its consequences.

4.1. Lens Model

In the context of a lens model, component 1 is a “straight arc” and component 5 is a “counterimage.” This gravitational lens image configuration is very common; it has been found in several clusters (see Surdej & Soucail 1993 for a review). The model for these systems is that of a source lying on or very close to a cusp in a caustic (a line of infinite magnification, e.g. Blandford & Narayan, 1992) in the source plane. Although

the magnification of a point source lying on the caustic is formally infinite, the maximum magnification of a real object is limited by its finite angular radius r . Under the gravitational lens hypothesis the total magnification of the source should be on the same order as the flux ratio of arc to counterimage, roughly 100 in this case. Gravitational lens models also predict that the axis ratio of the arc should be on the same order as the total magnification. The $0''.7$ length of the arc thus implies an observed width on the order of $0''.007$ (50 pc), or unresolved even in 11 ST images.

In the case of lensing dominated by mass at a single redshift, the gravitational lens mapping, which takes a two-dimensional angular position \vec{x} on the image plane (i.e. the position observed on the sky) to a two-dimensional angular position \vec{y} on the source plane (i.e. the position that would be observed if there was no lens), is a gradient mapping

$$\vec{y} = \vec{x} - \vec{\nabla}_{\vec{x}} \psi(\vec{x}), \quad (1)$$

where $\vec{\nabla}_{\vec{x}}$ is the two-dimensional gradient operator with respect to angular image-plane position \vec{x} , and $\psi(\vec{x})$ is a scaled, projected, two-dimensional gravitational potential. The potential is related to the angular surface density Σ (mass per unit solid angle)

$$\Sigma(\vec{x}) = \frac{c^2}{8\pi G} \frac{D_d D_s}{D_{ds}} \nabla_{\vec{x}}^2 \psi(\vec{x}), \quad (2)$$

where D_d , D_s , and D_{ds} are angular diameter distances from observer to lens (deflector), observer to source, and lens to source, and $\nabla_{\vec{x}}^2$ is the two-dimensional Laplacian operator,

The lens models which follow assume that the potential ψ can be approximated with a quasi-isothermal sphere with ellipticity (see, e.g. Kochanek, 1991), i.e.,

$$\psi(\vec{x}) = b \sqrt{s^2 + r^2} [1 - \gamma \cos 2(\theta - \theta_\gamma)], \quad (3)$$

where $\vec{x} = (r, \theta)$ is the position of the point in question relative to the center of the mass distribution, b is the asymptotic critical radius (the radius of the Einstein ring), roughly

the angular radius of the circle of images ($\sim 1''$ in this system because that is the angular separation of arc and lens), γ is an ellipticity parameter, θ_γ is the position angle of the major axis, and s is a core radius. The results do not depend strongly on the core radius s , so it is assumed to be zero. The critical radius b can be related to a one-dimensional velocity dispersion for the lens by

$$\sigma_v^2 = \frac{c^2 D_s}{4\pi D_{ds}} b, \quad (4)$$

although this depends on the assumption of isothermality. More secure is the mass M inside the “circle of images” (in this case a circle of angular radius b around component 2),

$$M = \frac{c^2 D_d D_s}{4 G D_{ds}} b^2. \quad (5)$$

The mass M and the inferred luminosity L of the lens can be used to compute a mass-to-light ratio as well. The inferred physical properties of the lens depend strongly on lens and source redshifts and weakly on world model. In this system the lens redshift is unknown, so σ_v , M , and M/L are given as a function of lens redshift for model 2 (see below) in Figure 4. Further discussion of figure 4 is deferred until section 4.3..

Model parameters b , γ , and θ_γ were varied to minimize (in a least-squares sense) the scatter in the source plane positions corresponding to the brightest 55 pixels in the deconvolved arc (figure 1 (d)) and the brightest pixel in the counterimage. In order to perform the minimization correctly given that the pixel locations are measured on the image plane, the Jacobian of the lens mapping was used to transform uniform image-plane uncertainties into correctly-weighted source-plane uncertainties.

The best-fit model parameters are given in Table 2. The inferred intrinsic source radius which makes the arc-counterimage flux ratio 100 is on the order of $0''.005$ (40 pc). The model makes the assumption that component 3 is a singular isothermal sphere ($\gamma = s = 0$) at the same redshift as component 2 and with critical radius $b_3 = 0''.6$, the expected value

under the assumption that components 2 and 3 have the same mass-to-light ratio. A simpler model which assumes that the potential is entirely due to an elliptical shaped mass centered on component 2 was also considered. The two-component model was adopted because the intrinsic ellipticity of the potential in this model is smaller than in the simpler model, in better agreement with the observed ellipticity in component 2. This is because the external mass of component 3 has a tidal effect which replaces some of the ellipticity in the primary lens. In both models the predicted orientation of the lens is consistent with the observed orientation of component 2. Figure 1(f) shows the density and potential contours for the adopted model, as well as the critical curve and the image morphology for a circular source of radius $0''.005$, smoothed to a FWHM of $0''.02$, and with the counterimage brightness enhanced for visibility. Figure 1 (e) shows the model image morphology convolved with the synthetic PSF discussed in section 3., and should be compared to figure 1 (c).

The image configuration in the lens model is that of a triple image or straight, arc (plus counterimage). Although parts of the source arc triply imaged in component 1, the source radius (~ 0.005 arcseconds) inferred from the flux ratio of components 1 and 5 is large enough that the three images merge into a single straight arc. We interpret the peak in the east half of the arc as corresponding to two images merging on the critical curve, while the peak in the west half corresponds to the third image. The triple structure may become more apparent in high-resolution images in other bandpasses if the flux at those wavelengths is produced by structures offset by $\sim 0''.02$ (160 pc) from those which produce the F814W flux, or have intrinsic size scales a factor of ~ 3 smaller.

The predicted total magnification of F814W emission from a uniform circular source as a function of source radius is shown in figure 5. The total magnification for a source radius of $0''.005$ is smaller than the flux ratio of component 1 to component 5, because component 5 is slightly demagnified. The dependence of the calculation of the total magnification in

the 11.5" image on the assumption of a circular source geometry for the F814W emission was investigated. Sources of the same total projected solid angle on the sky have the same total magnifications to within about 15% even if they are highly elliptical, no matter what their position angle. The magnification in the model scales as r^{-1} for very small sizes or separations from the caustic, and smoothly converts to $r^{-0.5}$ at larger radii, in agreement with Schneider, Ehlers & Falco (1992). The bump near $r \sim 0".5$ in figure 5 corresponds to the formation of a ring (see below). The magnification predicted by the model can be approximated to $\sim 20\%$ by $M = 3.9r^{-0.624}$ for the range $0".001 < r < 1"$ (S to S000 pc).

Different distributions for the narrow line and UV and optical continuum regions, and the likelihood of substantial reddening (Elston et al. 1994), can therefore account for the substantially different appearance of FSC 10214+4724 at different wavelengths noted by Matthews et al. (1994), and in particular for the larger extent of the K -band arc seen by Matthews et al. and Graham and Liu (1995) than the arc seen in the *HST* image. The 140° extent of the K -band arc corresponds to a source with $0".25$ (2 kpc) radius. If the source radius is increased to $\sim 0".5$ it is imaged into an elliptical ($e \sim 0.4$) ring connecting components 1 and 5. The position angle of this ring is perpendicular to that of component 2, and is offset from being perfectly centered on component 2 by $\sim 0".4$ in the direction of component 1. The excess light near 1".4 pixels noted in section 3.2. maybe the UV (rest frame) counterpart of the more extended arc seen in the K images. Note that Matthews et al. find the $H\alpha$ emission to be extended in an east-west direction by $\sim 0".5$, suggesting that the narrow line region is largely coincident with the UV continuum which dominates the 11'814W image.

4.2. Bolometric Luminosity of FSC10214+4724

FSC10214+4724 has an apparent luminosity of $L_{app} = 5 \times 10^{14} L_{\odot}$ (Rowan-Robinson et al. 1993), making it among the most luminous known objects in the Universe. The vast majority of this luminosity, $\sim 99\%$, is observed in the infrared (Rowan-Robinson et al. 1991, 1993). There is strong evidence that the UV source is a quasar (FWHM of C IV] $\sim 10,000 < 111 \text{ S}^{-1}$ in polarized light, Miller 1995) enshrouded in dust ($110/11.8 \geq 20$ implying $A_V > 5.5$, Elston et al. 1994), and that the quasar's luminosity is absorbed in the dust shell and reradiated in the infrared (Rowan-Robinson 1993). This implies that the size of the infrared emitting region is substantially larger than the optical/UV emitting region.

If FSC10214+4724 is magnified by a gravitational lens, the intrinsic source luminosity is less than the apparent luminosity. However, if the infrared source is larger than the optical/UV source, the magnification of the infrared source is less than the magnification measured from the *HST* image. The magnification of the infrared source can be estimated by assuming that the infrared source can be approximated as an optically thick blackbody. This assumption corresponds to making the infrared source as small as possible, and hence the magnification of the infrared radiation as large as possible. In this case, because of the assumption that the emitted infrared energy distribution is independent of distance from the central heating source, the magnification is independent of wavelength. The temperature of the dust is assumed to be $T \sim 140 \text{ K}$. At this temperature the emission peaks at a rest wavelength of $18 \mu\text{m}$, corresponding to the observed emission that peaks at $60 \mu\text{m}$.

With this temperature, the apparent luminosity L_{app} and intrinsic luminosity L_{int} can be written as

$$L_{app} = M(R)L_{int} = M(R) \times 4\pi R^2 \sigma T^4 \quad (6)$$

where R is the physical radius of the source, $M(R)$ is the magnification from figure 5 for a uniform disk of radius R , and T is the blackbody temperature determined by the wavelength of peak emission. Solving this equation for R gives a radius of 130 pc ($0''.017$),

and $M(R) = 42$, so that the intrinsic luminosity of FSC 10214-4724 is $1.2 \times 10^{13} L_{\odot}$. A slightly larger source size and lower magnification is derived if the temperature T is assumed to be 115K, the color temperature determined by the observed flux densities at 60 and $450 \mu\text{m}$ and corrected for redshift. Then the radius of the infrared source is 240 pc ($0''.03$), the magnification is 29, and the intrinsic luminosity is $1.7 \times 10^{13} L_{\odot}$.

The expected arc length is $\sim 2RM(R)$, or $1''.7$ in the $T = 115\text{K}$ case, and $1''.4$ for $T = 140\text{K}$. From VI, A-A configuration observations at 8.4 GHz with $0''.25$ resolution, Lawrence et al. (1993) found a $0''.6$ (east-west) by $1''.3$ source. The similarity of this structure to the arc in the *HST* image suggests a continuum radio source radius closer to the $0''.005$ (40 pc) estimated for the optical/UV source than to the minimum far-infrared size just calculated. Condon et al. (1991) find that the radio source size for nearby IRAS galaxies with far infrared luminosities $> 10^{12} L_{\odot}$ is typically ~ 100 pc (and for Mrk 231, the most luminous of the sample, $\lesssim 1$ pc), smaller than the minimum blackbody size for far infrared emission from these galaxies. For their sample Condon et al. find $\langle q \rangle = 2.34$, where q is the logarithm of the ratio of far infrared (60-100 μm) to 1.49 GHz flux. For FSC 10214-4724, extrapolating the Lawrence et al. (1993) observed radio flux to 0.45 GHz (the observed frequency for emitted 1.49 GHz) yields 3.5 mJy, and interpolating to the rest frame wavelengths for 60 and 100 μm and using Condon et al.'s definition gives $q = 1.91$. If the radio magnification is 100, and the far infrared magnification is 30, then the intrinsic $q = 2.39$. Therefore the radio morphology and flux measured by Lawrence et al. are quite consistent with the above estimate for the bolometric luminosity.

The $0''.6$ extent of the radio morphology is also consistent with a much smaller radio continuum source size, although the value of q would then be significantly larger than observed for local luminous IRAS galaxies. It would be interesting (albeit quite challenging) to see whether the very high angular resolution possible with VLBI observations revealed

the triple structure in the arc discussed above.

The size of the infrared source determined under the assumption of optically thick emission is a plausible lower limit to the physical source size. Alternatively, the magnification can be estimated based on the models of Phinney (1991) of infrared emission from dusty, warped disks illuminated by a central quasar. The physical size of the source required to obtain a self-consistent solution for the intrinsic luminosity is quite large. For the region emitting at $150\mu\text{m}$ ($450\mu\text{m}$ observed), the source radius would be $\sim 10''$, much larger than the observed size of the CO source (e.g. Scoville et al. 1995). Thus we consider such a model less consistent with the observations than the optically thick models.

The reduction in the intrinsic luminosity of FSC10214+4724 to $\sim 2 \times 10^{13} L_{\odot}$ implied by the lens model of the source brings it into the luminosity range of previously studied infrared luminous AGN. The IRAS source FSC15307+3252 at a redshift of $z=0.93$ has a luminosity of $4 \times 10^{13} L_{\odot}$, while the IRAS source PSC09104+4109 has a luminosity of $2 \times 10^{13} L_{\odot}$ for our assumed cosmology (Cutri et al. 1994). There is no known evidence from high resolution imaging (Soifer et al. 1994, Hutching and Neff 1988, Soifer et al. 1995 in preparation) that either of these sources is a gravitational lens, so the apparent luminosity is presumably the intrinsic luminosity in these cases. Thus, based on its bolometric luminosity, FSC10214+4724 is most likely a source similar to these. The reduction in intrinsic luminosity reduces the necessary dust mass associated with the source (Rowan-Robinson et al. 1993) by the same magnification factor, into the range $M_{dust} \sim 1 - 3 \times 10^7 M_{\odot}$, which is consistent with the estimates of the gas mass based on the dynamical mass determinations from the CO observations (Scoville et al. 1995).

4.3. Properties of Component 2

No conclusive measurement of the redshift for component 2 has yet been made, to our knowledge, although values of 0.42 (Close et al. 1995) and 0.90 (Serjeant et al. 1995) have been suggested based on possible continuum breaks in the spectrum of component 2, while Miller (1995) finds Mg lines in absorption at $z = 1.32$ (and possibly $z = 0.89$) in the spectrum of component 1. In the Appendix we provide three estimates of the redshift for component 2 (two of which are closely related). All three estimates are consistent with $z \sim 0.9$, and we adopt this value as our best overall estimate of the redshift. Note the SED and $R_c - < \mu_B >_\epsilon$ estimates do not assume component 2 is a lens, only that it is an elliptical galaxy, and therefore give additional support to the lensing hypothesis by placing component 2 at an intervening redshift relative to FSC10214+4724.

The velocity dispersion σ_v , mass M , and mass-to-light ratio (M/L) predicted for the lens are shown in figure 4 as a function of lens redshift. Adopting $z = 0.9$ yields $(M/L)_B = 8M_\odot/L_\odot$ (vs. the observed average of 6, van der Marel 1991), $\sigma_v = 270$ km s⁻¹, and $M = 3.8 \times 10^{11} M_\odot$ (thus $L_B = 5 \times 10^{10} L_\odot$). These values are for a radius of 0".85: using figure 3 the total blue luminosity is then $L_B = 1.4 \times 10^{11} L_\odot$ or $\sim 4L^*$ (Binggeli, Sandage and Tammann 1988). These values are independent of evolutionary model because F814W samples rest frame B at $z = 0.9$. Thus for the redshift estimate $z = 0.9$ the lensing model predicts properties typical of present day elliptical galaxies, except that the galaxy is unusually luminous. The probability of a large lensing galaxy is greater than the galaxy luminosity function alone implies, however, because the cross section for gravitational lensing is proportional to mass.

4.4. The Parent Population of IRAS FSC10214+4724

Analysis of statistically complete samples of radio galaxies suggests that the lensing rate (i.e. probability that a given radio galaxy is lensed) is on the order of 1/500

(Miralda-Escudé & Lehar, 1992; CLASS survey, private communication). Given that a source is lensed, the probability of getting total magnification M is on the order of M^{-2} (e.g. Schneider, Ehlers & Falco, 1992). The estimated total magnification ~ 30 for the IRAS flux from section 4.2. corresponds to a likelihood of $\sim 10^{-3}$. The existence of a single lensed object in the surveyed area (0.2 sr - Rowan-Robinson 1991) with magnification 30 should, according to these probabilities, represent an underlying population of ~ 800 compact, $60\mu\text{m}$ -luminous objects per square degree (or >40 per square degree at 95% confidence) which are either not lensed or lensed with much lower magnification (and hence are not in the FSS catalog). These sources will have observed magnitude $r \sim 25$ mag if they are like IRAS FSC10214+4724 and their IR fluxes will be of order 3 mJy at $25\mu\text{m}$ and 7 mJy at $60\mu\text{m}$. At these flux levels, models of the IR galaxy population with strong luminosity evolution (Hacking and Soifer, 1991) predict a few hundred sources per square degree, in agreement with this estimate. Of course this is only an order of magnitude estimate because it depends on extrapolation from a single serendipitously discovered object, and on the relative redshift distributions of IR-luminous and radio galaxies. optical field galaxy redshift surveys now underway with the Keck Telescope are approaching this depth (J. Cohen, private communication; UCDEEP collaboration, private communication), and IR imaging surveys to well beyond these levels are envisioned with ISO, WIRE, and SIRTF, so this very uncertain prediction may be testable in the near future.

5. Summary

We have obtained a O.Spin image of the $z=2.286$ IRAS source FSC10214+4724 with the *HST* WFPC2 Planetary Camera, with $0''.1$ resolution and high signal to noise. We find the following:

- 1) The source appears as an unresolved ($< 0''.06$) arc $0''.7$ long, with significant

substructure along its length. The arc is centered on a galaxy $1''.18$ to the north (component 2), and a faint unresolved component (component 5) is clearly detected $0''.43$ north of component 2. Two other galaxies (components 3 and 4) are evident within a few arcseconds of the IRAS source. This morphological configuration is characteristic of a gravitationally lensed system, in which the arc and component 5 are images of a single background source produced by the potential of the foreground component 2.

2) The surface brightness profile of component 2 is well matched by a de Vaucouleurs profile, characteristic of an elliptical galaxy with an effective radius of $1''.27$. There is evidence for excess emission above the de Vaucouleurs profile near the radius of the arc.

3) The flux ratio of the arc to the component 5 is ~ 100 , implying magnification in the *HST* image of the background source by roughly this amount.

4) A detailed lensing model which reproduces the observed morphology and relative flux of the arc and counterimage correctly predicts the position angle for component 2. Better agreement is found with the observed ellipticity of component 2 if component 3 is included in the lensing potential. The model predicts reasonable values for the mass and velocity dispersion of component 2.

5) If component 2 is an elliptical galaxy, its spectral energy distribution is inconsistent with it being at $z = 2.286$, and $z = 0.9$ is preferred. The surface brightness profile of component 2 implies a redshift between 0.6 and 1.2. From the lensing model, for $z \sim 0.9$, the central mass-to-light ratio for component 2 is $(M/L)_B = 8M_{\odot}/L_{\odot}$, the velocity dispersion $\sigma_v = 270 \text{ km s}^{-1}$, and the total blue luminosity $L_B = 1.4 \times 10^{11} L_{\odot} \sim 4L^*$.

6) The model predicts an intrinsic radius of $\sim 0''.005$ (40 pc) for the background source. Triple structure in the arc is obscured by this source size, but may become apparent at high resolution in other bandpasses. The larger size of the arc observed at *K* implies an intrinsic

source radius of $0''.25$ in the corresponding emitting bandpass. A source of radius $> 0''.5$ would produce a ring of emission connecting the arc and component 5. This may account for the excess emission seen in the surface brightness profile of component 2. The $H\alpha$ and radio continuum morphologies appear similar to that of the $0.8\mu\text{m}$ arc, implying a similar source size for the narrow line, UV continuum, and radio continuum emission.

i) The minimum source size for an optically thick blackbody source producing the bulk of the bolometric luminosity is $\sim 0''.03$ (240 pc), implying a bolometric magnification of ~ 30 . The background lensed source then has an intrinsic luminosity $\sim 2 \times 10^{13} L_{\odot}$. Thus IRAS FSC10214+4724 is not the most luminous object in the Universe, but it remains among the most luminous in the IRAS catalog.

S) The expected incidence of 30-fold gravitational magnification is low enough to suggest that FSC10214+4724 represents an underlying population of ~ 500 compact, objects per square degree with optical magnitude $r \sim 25$ mag and $F_{60\mu\text{m}} \sim 7$ mJy.

We thank Mark Dickinson for help with the $R_e - \langle \mu_B \rangle_e$ technique for estimating z and in particular for supplying the Sandage & Perlmutter data in electronic form, Adam Stanford for calculating K-corrections and general assistance with STSDAS, and Roger Blandford for help with lens modelling. This research was supported by NASA through a grant awarded by STScI, which is operated by AURA under NASA contract, NAS 5-2655,5. Portions of the research described in this paper were carried out by the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA.

A. Estimates of the Redshift for Component 2

Here we use the spectral energy distribution and surface brightness profile of component 2 to estimate its redshift.

A.1. Spectral Energy Distribution

Figure 6 combines R and H data from Elston et al. (1994), J and K from Matthews et al. (1994), and F814W data from the present work for components 2, 3 and 4, normalized at K . (Note that component 1 in the Elston et al. terminology is our component 2.) Because the angular resolution of the three data sets ranges from $0''.1$ – $1''.5$, the combined spectral energy distribution (SED) is somewhat uncertain.

Given the good agreement of the surface brightness profile for component 2 with a de Vaucouleurs law (figure 3) it is reasonable to assume that this component is an elliptical galaxy. For comparison, the unevolved spectrum of a standard Bruzual and Charlot (1993) elliptical galaxy model at an age of 13 Gyr and redshifts of 0.42, 0.90, 1.32, and 2.286 is plotted in figure 6(a); the corresponding passively evolving model with ages of 7.75, 5, 3.75, and 2.2 Gyr at these redshifts (the ages are consistent with a present age of 13 Gyr with the assumed cosmology) is plotted in figure 6(b). All models were normalized to the K flux in the SED for component 2. Clearly the $z = 2.286$ models fail to match the observed SED for component 2, while the $z = 0.9$ models provide surprisingly good fits to the observations. This is fairly strong evidence that component 2 is in fact a foreground elliptical: it is too blue to be an elliptical galaxy at the redshift of FSC10214+4724.

The r data point for component 3 is anomalously bright, while the rest of its SED is somewhat redder than component 2. This might be due to a combination of reddening and star formation associated with the tidal interaction suggested in section 3.1., placing

component 3 at the same redshift as component 2, as Model 2 in section 4.1, assumes. Alternatively, the $z = 1.32$ models appear to fit component 3 at least as well as the $z = 0.9$ models. Without the constraint of an elliptical surface brightness profile, it is much more difficult to assign a unique redshift based on the SED. Component 4 is significantly bluer than components 2 and 3, consistent with what appears to be a later type morphology.

A.2. Fundamental Plane Relations

Since the surface brightness profile of component 2 strongly suggests it is an elliptical galaxy (section 3.2, and figure 3), it is possible to make further use of the surface brightness profile to estimate the redshift of component 2 using the fundamental plane relations for ellipticals (Kormendy and Djorgovski 1989). Using $r_e = 1''.274$ and $m_{814} = 21.04$ within r_e from figure 3, the present day equivalent blue surface brightness of component 2 was calculated by correcting for $(1+z)^4$ surface brightness dimming, redshift K-correction, and luminosity evolution. Figure 7 shows R_e (in kpc) and $\langle \mu_B \rangle_e$ (the average blue surface brightness within R_e) as a function of the assumed redshift for component 2, overlaid on the data for present day ellipticals from Sandage and Perelmutter (1990). Luminosity and K-corrections are shown for both a non-evolving and passively evolving elliptical model spectrum from Bruzual and Charlot (1993). Redshifts near zero, or in the range 0.6 to 1.2 can be accommodated. From figure 4, the lens model predicts a central mass-to-light ratio > 100 for $z < 0.2$, arguing against, low values. We consider the passively evolving model more realistic, leading to an estimate of $z = 1.0 \pm 0.2$. This estimate is independent of H_0 because the present day data scale in the same way as the calculated values. The estimate is driven almost entirely by the $(1+z)^4$ dependence of surface brightness on redshift, and is relatively independent of q_0 because the latter primarily affects angular size, which is nearly orthogonal to redshift in the region of interest in figure 7. The main uncertainty is due to

luminosity evolution (some of which arises from the dependence of timescales on q_o) and to the scatter in surface brightness among giant elliptical galaxies.

Much of this scatter is correlated with the **velocity** dispersion, and if σ is known the $D_n - \sigma$ relation (Lynden-Bell et al. 1988) can be used to measure the angular diameter distance for giant elliptical galaxies. Here D_n is defined as the angular diameter of the circle within which the integrated rest frame blue surface brightness is $20.75 \text{ mag arcsec}^{-2}$ after correction for luminosity evolution and $(1+z)^4$ surface brightness dimming. The value of σ is found from the lens model as plotted in figure 4(a). An advantage of D_n over R_e is that D_n is defined at a high surface brightness level and is therefore smaller than R_e and immune to uncertainties about emission at the arc radius, as well as being less sensitive to uncertainties in sky subtraction. However the technique is sensitive to q_o because it is essentially an angular diameter distance. The redshift estimates from this approach range from 0.75 for the case $q_o = 0$ and no evolution, to 1.15 for $q_o = 0.5$ and passive evolution, independent of l_1 .

| Component | m_{814} | $\Delta\alpha''$ | $\Delta\delta''$ | Comment |
|-----------|-----------|------------------|------------------|-----------------------------|
| 1 | 20.44 | 0.10 | -1.16 | includes faint extensions |
| 2 | 21.41 | 0 | 0 | inside $r = 0''.85$ |
| 2 | 20.3 | | | total |
| 3 | 23.16 | 1.03 | 1.93 | inside $r = 0''.5$ |
| 3 | 22.98 | | | including component to east |
| 4 | 23.58 | 3.42 | 1.79 | |
| 5 | 25.5 | 0.03 | 0.43 | |
| Star 11 | 24.54 | -6.59 | -6.20 | |

Table J : Photometry Of objects in *HST* F814W image of IRAS FSC10214+4724.

Note. - Components are identified in figure 1 (a) and (b). Positions are with respect to the center of component 2, which is approximately $\alpha = 1011241113''.56$, $\delta = 47^{\circ}09'10''.8$, (J2000) in the *HST* guide star catalog frame. For component 1 the position is for the peak brightness.

| Parameter | Model Value | observed | Comments |
|-----------------|-------------|----------------|---|
| b | 0.82 | | arcseconds |
| γ | 0.12 | | defined by Eqn. (3) |
| c | 0.30 | 0.16 ± 0.1 | $1 - b/a$ |
| θ_γ | -11 | 3 ± 15 | degrees, N of E |
| b_3 | 0.60 | | arcseconds |
| source radius | 0.0051 | | arcseconds, for magnification ratio 100 |

Table 2: Lens Model Parameters.

Note. The model includes component 3 in the lensing potential. The ellipticity c is the conventional value defined by one minus the ratio of semiminor to semimajor axis, and differs from the model ellipticity parameter γ which is defined by Eqn. (3). The “source radius” is the angular radius at which, for a circular source, the arc-counterimage magnification ratio is 100. Other symbols are explained in the text. World model $q_0 = 0.5$ is assumed. Changing world models only changes the numbers by $\sim 10\%$.

REFERENCES

- Binggeli, B., Sandage, A., & Tammann, G. A. 1988, *ARA&A*, 26, 509
- Blandford, R. D. & Narayan, R. 1992, *ARA&A* 30, 311
- Broadhurst, T. J. & Lehar, J. 1995, *ApJ*, in press
- Brown, R. L., & Vanden Bout, P. A. 1991, *AJ*, 102, 1956
- Bruzual, G., & Charlot, S. 1993, *ApJ*, 405, 535
- Clements, D. L., Rowan-Robinson, M., Lawrence, A., Broadhurst, T., & McMahon, R. 1992, *MNRAS*, 256, 35P
- Close, L. M., Hall, P. B., Liu, C. T., & Hege, E. K. 1995, *ApJ*, submitted
- Condon, J. J., Huang, Z.-P., Yin, Q. F., and Thuan, T. X. 1991, *ApJ*, 378, 65
- Cutri, R. M., Huchra, J. P., Low, F. J., Brown, R. L. & Vanden Bout, P. A. 1994, *ApJ*, 424, L65
- Davies, R. L., Efsthathiou, G., Fall, S. M., Illingworth, G., & Schechter, P. L. 1983, *ApJ*, 266, 41
- Downes, D., & Radford, S. J. E., Greve, A., Thum, C., Solomon, P. M., & Wink, J. E. 1992, *ApJ*, 398, 1,25
- Elston, R., McCarthy, P. J., Eisenhardt, P., Dickinson, M., Spinrad, H., Jannuzi, J. P., & Maloney, P. 1994, *AJ*, 107, 910
- Fisher, D., Illingworth, G., & Franx M. 1995, *ApJ*, 438, 539
- Graham, J. R. & Liu, M. C. 1995, *ApJ*, in press
- Hacking, P. L. & Soifer, B. T. 1991, *ApJ*, 367, L49
- Hutchings, J. B. & Neff, S. G. 1988, *AJ*, 96, 1575.

- Kochanek, C. S. 1991, ApJ 373, 354
- Kormendy, J., & Jorgovski, S. 1989, ARA&A, 27, 235
- Lawrence, A., et al. 1993, MNRAS, 260, 28
- Lynden-Bell, D., Faber, S. M., Burstein, D., Davies, R. L., Dressler, A., Terlevich, R. J., & Wegner, G. 1988, ApJ, 326, 19
- Matthews, K., et al. 1994, ApJ, 420, L13
- Miller, J. S. 1995, BAAS, 27, 869
- Miralda-Escudé J. & Lehar J. 1992, MNRAS 259, 311
- Phinney, E.S. 1989, *Theory of Accretion Disks* W.Duschl, F.Meyer and J. Frank, eds (Dordrecht: Kluwer Academic Publishers) p 111
- Rowan-Robinson, M. et al. 1991. *Nature*, 351, 719.
- Rowan-Robinson, M. et al. 1993, MNRAS, 261, 513.
- Sandage, A., & Perlmutter, J. 1990, ApJ, 361, 1
- Schneider, P., Ehlers, J., & Falco, E. E. 1992, *Gravitational lenses*, Springer-Verlag, New York.
- Serjeant, S., Lacy, N.J., Rawlings, S., King, I., & Clements, D. L. 1995, MNRAS, in press
- Scoville, N. Z., Yun, M. S., Brown, R. L. & Vanden Bout, P. A. 1995, ApJ, in press
- Soifer, B. T., Neugebauer, G., Matthews, K., & Armus, L. 1994, ApJ, 433, L69.
- Solomon, P. M., Downes, D., & Radford, S. J. E. 1992, ApJ, 398, L29
- Surdej J. & Soucail G. 1993, in *Gravitational Lenses in the Universe*, ed. J. Surdej et al., Proc. 31st Liège Int. Astroph. Conf., pp. 205-216.
- Trentham, N. 1995, MNRAS, submitted
- Tsuboi, H., & Nakai, N. 1992, PASJ, 44, 1241

Vail der Mareel, 1{. 1'. 1995,MNRAS, 253, 710

Figure Captions

Figure 1: Montage of HST Planetary Camera (PC) imaging of IRAS FSC10214+4724 in F814W. Panels (a), (b), and (c) show the image at progressively finer scales, as indicated by the axes which are labelled in arcseconds relative to component 2 (see panel (b)). Panels (d), (e), and (f) have the same scale and center as panel (c). Panel (d) is a partially deconvolved version of the data with a factor of four subsampling. Panel (e) shows the observed image configuration given the lens model and a uniform circular disk source of radius 0.0055 arcseconds. The image configuration has been convolved with the synthetic *HST* PSF and binned into PC pixels to allow direct comparison with panel (c). Panel (f) shows further details of the lens model: as in panel (c) the pixel size is four times smaller, and the model image has been lightly smoothed to a FWHM of $0''.02$. Lines indicate contours of mass (dotted), potential (dashed), and the critical curve (solid line) for the model. The grey levels in panel (f) range linearly from zero (white) to the peak value in the arc (black), but have been enhanced (in panel (f) only) by a factor of seven at the counterimage location. The grey levels in panels (a) - (c) range linearly from 0.5% to 5% of the peak brightness in component 1 ($\mu_{814} = 17.6 \text{ mag arcsec}^{-2}$ in panels (a) (c), $\mu_{814} = 13.6 \text{ mag arcsec}^{-2}$ in panel (d)). Contour levels in panels (c) (e) are at 25, .50, 75 and 90% of this peak brightness for component 1, or at 25, 50, 75 and 90% of the peak brightness for component 2 ($\mu_{814} = 19.4 \text{ mag arcsec}^{-2}$ in panel (c), $\mu_{814} = 15.6 \text{ mag arcsec}^{-2}$ in panel (d)), as appropriate. North is $37.1''$ counterclockwise from vertical in all panels, with east 90° counterclockwise from north, as shown in panels (a) and (b).

Figure 2: Radial profiles for objects identified in figures 1 (a) and (b). Component 1 appears unresolved relative to stars A and 11. For component 1 the equivalent radial profile is plotted, as discussed in section 3.2. The synthetic PSF was used to generate the deconvolution shown in Figure 1 (e). The vertical scales for the profiles were normalized

at the smallest radius available, except for star A, whose *core* is saturated in our image, and which was normalized to the synthetic PSF at the first radius which was not saturated. The data points plotted for negative radii are identical to those for positive radii except for component 1.

Figure 3: The surface brightness of component 2 is plotted as a function of radius, together with a deVaucouleurs profile with $r_e = 1''.27$ (solid line) and an exponential profile with $r_d = 0''.32$ (dotted line). Magnitude as a function of aperture radius is shown by the open circles and the righthand scale. The magnitude appears to converge near $m_{814} = 20.3$ mag, yielding a consistent value for r_e .

Figure 4: Predictions from the lens model (see section 4.1. and Table 2) for the mass inside the critical radius (panel b) and the velocity dispersion (panel a) of the lens as a function of its redshift. The bottom panel (c) shows the mass to luminosity ratio using the *HST* F814W flux inside $0''.85$ radius, K-corrected to rest frame *B* using unevolving (solid curve) and passively evolving (dotted curve) elliptical model spectra from Bruzual & Charlot (1993). The dashed vertical line indicates the best estimate for the redshift of component 2 as discussed in the Appendix. Horizontal dashed lines show the normally observed range of $(M_{\odot}/L_{\odot})_B$ and σ_v from Fisher, Illingworth, and Franx (1995), van der Marck (1991), and Davies et al. (1983). The value $M^* = 2 \times 10^{12} M_{\odot}$ is from van der Marel's mean $(M_{\odot}/L_{\odot})_B = 6$ and $L_B^* = -21$ from Binggeli, Sandage and Tammann (1988). The values shown are for the assumed cosmology ($H_0 = 50$ km S⁻¹, $q_0 = 0.5$), and are relatively insensitive to q_0 . The mass scales as h^{-1} and the mass to luminosity ratio as h (for the unevolving K-correction), while the velocity dispersion is independent of h , where h is the Hubble constant in units of 100.

Figure 5: Predictions from the lens model for the total magnification M of the background source flux, assuming a uniformly illuminated source of radius r (arcseconds,

shown on the bottom axis) or R (parsecs, shown on the top axis). The dotted line shows a power law approximation to the predicted magnification.

Figure 6: Spectral energy distributions for components identified in figure 1(b), with unevolving (panel a) and passively evolving (panel b) model elliptical spectra from Bruzual & Charlot (1993) shown for comparison. The component 2 data fit the $z = 0.9$ models well. The flux scale is correct for component 2; for component 3 it should be reduced by a factor of 2.3 and for component 4 by a factor of 9. The data are derived from Elston et al. (1991), Matthews et al. (1994), and this paper. See section A.1 for further details.

Figure 7: Estimate of the redshift for component 2 using the average blue surface brightness - effective radius relation for giant elliptical galaxies. The data are from Sandage and Perlmutter (1990). The dotted curve shows calculated $\langle \mu_B \rangle_r$ and R_e values with K-corrections from an unevolving Bruzual and Charlot (1993) elliptical model with $q_0 = 0$ and a uniform redshift interval of 0.05. Here $\langle \mu_B \rangle_r$ is the average blue surface brightness within the effective radius. The solid curve is for a passively evolving elliptical and $q_0 = 0.5$.







Fig. 1



Component 2 Radial Surface Brightness Profile




Lens Properties $H_0 = 50 \text{ km/s}$ $q_0 = 0.5$



Source Radius (pc, $H_0 = 50 \text{ km/s}$, $q_0 = 0.5$)

..... $M = 3.9 \text{ r(arcsec)}^{-0.624}$



Source Radius arcseconds

0.01 0.1 1

Total Magnification

No Evolution

Passive Evolution

