

**Time-Dependent Hydrodynamic Modeling of the
Global Structure of the Heliosphere**

P. C. Liewer, (Jet Propulsion Laboratory, California Institute
of Technology, Pasadena CA 91109; 818-354-6538)
S. Roy Karmesin, (California Institute of Technology,
Pasadena CA 91125)
J. U. Brackbill, (LANL, Los Alamos, NM 87545)

We present results from time-dependent two-dimensional hydrodynamic modeling of the global structure of the heliosphere. The solar wind plasma is assumed to flow out radially with a uniform supersonic velocity. Three cases are considered. First, we consider the case of a subsonic external local interstellar medium flow and compare results for the termination shock and stagnation point locations with the incompressible models of Parker[1] and Suess and Nerney[2]. Second, we consider the case of a supersonic external flow and compare with the results of Baranov[3]. Third, the solar wind velocity is allowed to vary in time and we study the effect of variable solar wind velocity on the location and speed of movement of the termination shock.

1. E. N. Parker, *Interplanetary Dynamical Processes*, (John Wiley and Sons, NY) 1963.
2 S. T. Suess and S. Nerney, J. Geophys. Res. 95, 6403 (1990).
3. V. B, Baranov and Yu. G. Malama, J. Geophys. Res. 98, 15,157 (1993).

* Supported by NASA/H eliospheric Physics

1. 1994 Spring Meeting

2. 011216771 (F. Liewer)

3. (a) **P. C. Liewer**
MS 169-506
JPL
4800 Oak Grove Dr.
Pasadena, CA 91109

(b) 818-354-6538

(c) 818-354-8895
9-i

(b) **2124 Heliopause and Termination**
2164 Solar Wind Plasma
2199 General

0 %

8. charge \$50 to VISA
4024-0046-5349-8556

9. C

10. Oral Presentation Requested

No