

## Experiments Along Coexistence near Tricriticality in 3He-4He Mixtures

Melora Larson(a), Vladimir Dotsenko(b), Norbert Mulders(b), Alfred Nash(a), John Panek(a, 1), and Ben Vollmayr-Lee(d)

(a) Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109

(b) Department of Physics and Astronomy, University of Delaware, Newark, DE 19716

(d) Department of Physics, Bucknell University, Lewisburg, PA 17837

The tricritical point in the phase diagram of 3He-4He mixtures offers unique opportunities to test our understanding of critical phenomena. Because  $D = 3$  is the marginal spatial dimension for tricriticality, the associated critical exponents are exact integer fractions. In addition, one expects to find logarithmic corrections. We present our recent results for the superfluid density, obtained from time-of-flight non-linear second sound measurements near the tricritical point and along the phase separation curve. We also report on our measurements of the phase separation curve near the tricritical point performed using inter-digital capacitor sensors on the top and bottom of our cell. With our sensors, we are able to measure the location of the phase separation curve on both sides of the tricritical point, as well as the superfluid transition line as it approaches the tricritical point. We also will show our measurements of film behavior both near the superfluid transition and near the tricritical point.

(1) Current address: Goddard Space Flight Center, Greenbelt, MD 20771