

# Playing “Hide-and-Seek” With Binary Stars:

Searching for companions and accretion  
disks in AGB stellar systems

Elin Deeb

Mentor: Dr. Raghvendra Sahai




Grant No.  
0952013 &  
0833353

S. D. BECHTEL, JR.  
FOUNDATION  
STEPHEN BECHTEL FUND




# Introduction



Planetary Nebulae  
(PN/PNe) come  
in a variety of  
shapes


# Introduction



- Why do stars evolve from spherical objects into asymmetric PNe?
- How does this impact the recycling of material back into the galaxy?
- Theories suggest companion stars may play a role (Sahai 2012)

# Asymptotic Giant Branch (AGB) Stars

- Phase prior to PN stage
- “Red Giant stars”
- Inert C/O core from He fusion
- Surface temperature has decreased
- Variable stars: luminosity changes periodically (~1 year cycle).




# Challenges Seeking Companions



- One exception: Mira is a binary system consisting of Mira A: an AGB star, and Mira B: a WD companion (Riemers & Cassatella 1985)
- White dwarf (WD) & main sequence (MS) stars are smaller and thus less luminous
- Variability of the AGB star prevents certain methods of detecting the companion from being viable (Sahai 2008)

# Blackbody Spectra



Hotter objects  
emit larger  
amounts of  
radiation as  
well as higher  
energy  
radiation

UV spectra were taken by GALEX of the following objects:

Name	Spectral Type (SIMBAD)	Variable Type
WPEG	M7e D	Mira
EP AQR	M8IIIv	SRb
R UMA	M3e-M9e D	Mira (O rich)
AA CAM	M5 S	Pulsating/Irr (O rich)
EY HYA	M7IIIa D	SRa
TZ HOR	M5III C	Pulsating Variable
V ERI	M5/M6IV C	SRb (O rich)
TW HOR	N0 C7	SRa (C rich)
VY UMA	C5III	SRb (C rich)
V HYA	N:C7,5	Mira (C rich)

# Process

- Determine UV emission lines that can be emitted by AGB stars
- Determine UV emission lines that have been found in AGB binaries, as well as other interacting or accreting systems
- Use GALEXSpec to reduce data
- Use IRAF to plot data and measure central wavelength, flux, fwhm, and wavelength range
- Calculate flux/noise


## Ultraviolet Observations in AGB Systems

AGB stars (although cool) emit SOME UV light (possibly chromospheric emission):

- Mg h & k lines (centered at 2800 Å)
- Fe II lines (various) (Judge & Jordan 1991)
- C II features centered around 2325 Å  
(Johnson & Luttermoser 1987; Stencel 1982)

AGB stars emit relatively little flux at wavelengths less than 2800 Å due to their temperatures.

# EY Hya—our “poster child”



# EY Hya Line Identifications

$\lambda$ peak (Å)	$\lambda$ average (Å)	wavelength range	Flux (ergs cm <sup>-2</sup> s <sup>-1</sup> Å <sup>-1</sup> )	gfwhm	Identity	Max Flux/Noise
1454.0	1451.0	1443-1461	6.52E-015	3.960	N III?	3.88
1467.0	multiplet?	1461-1482	7.75E-015	8.024	C I or C II multiplet?	3.935
1485.5/1492.5	1489.0	1482-1496	4.79E-015	3.099	N IV]	4.10/4.30
1500.0	1499.5	1496-1517	1.66E-015	2.174	N III?	4.33
1531.0	1531.0	1528-1538	1.91E-015	1.934	C II multiplet?	5.06
1545.0	1545.0	1538-1550	1.86E-015	3.637	C IV	5.44
1587.0	1590.0	1584-1598	2.94E-015	4.022	unknown	4.59
1629/1643	1636.0	1625-1647	1.22E-014	3.037	He II	4.71/4.08
1660/1666	1660.5	1647-1674	1.99E-014	10.049	O III	3.13
1720.0	1719.0	1713-1724	3.75E-015	3.163	N IV	2.77
1727/1734	1731.0	1724-1737	4.70E-015	3.330	N III?	2.47/2.56
1741.0	1746	<b>1738-1755</b>	<b>6.32E-015</b>	<b>4.79</b>	N III]	3.09
1758.5/1765.5	1762	<b>1751-1773</b>	3.54E-014	21.149	C II	3.48
2157.0	2151.0	2136-2175	8.43E-015	33.110	Fe I or C II?	5.04
2332.0	2333.0	2312-2353	1.30E-014	31.010	C II	5.78
2452	2455	2437-2476	7.32E-015	31.88	Fe II	5.53
2620	2629	2606-2648	2.35E-014	58.12	Fe II	5.63
2728, 2738.5, 2752.5, 2756	multiplet	2709-2770	4.73E-014	58.23	Fe II	4.26/4.16
2798.0	~2800	2781-2818	7.09E-014	32.850	Mg II h and k	4.66

# The Resolution Challenge...


- The low resolution of the GALEX grism made it challenging to identify lines...


# The Resolution Challenge...

Example:



“Face on Mars”

Viking 1






# The Resolution Challenge...



# W Peg

Wait...what spectrum?


Phase affects the UV output of AGB stars (Luttermoser 2000)




## EP Aqr

Asymmetric double line profiles seen in CO lines (Kerschbaum & Oloffson 1999). C II line shows similar structure.

Common in O-rich AGBs. Suggested to be caused by a multiple shell structure with differential mass loss rates



# V Hya



# Challenges Faced

- The low resolution certainly affected the observations and analysis
- FUV observations are only available for EY Hya as there was an instrument failure
- The emissions of variables can vary by phase, which may impact the observations and analysis (Luttermoser 2000).
- Some spectra had very odd shapes, likely due to winds (Shore 1993).

# Summary

- FUV observations are critical to identifying companions. NUV features are theorized to be primarily AGB chromospheric features (Riemers & Cassatella 1985; Judge and Jordan 1991; Johnson & Luttermoser 1987).
- EY Hya is the only object with FUV data. Due to the low resolution, the findings thus far only weakly suggest an accretion disk (and thus a companion) may be present
- Other researchers indicate V Hya likely has a companion (Evans 2009), but our data is inconclusive.
- Phase, line shape, and chromospheric dynamics are important to consider

# Next Steps

- A proposal is being written to obtain FUV via the Cosmic Origins Spectrograph (COS) on board the Hubble Space Telescope. The lowest resolution still exceeds the resolution of GALEX.
- Additional NUV observations may be made as well due to the resolution challenges faced
- X-ray observations may also be considered down the road.



# Acknowledgements

The author would like to thank Dr. Raghvendra Sahai for his great mentorship as well as Petra Kneissl-Milanian for her role in the JPL internship program. The author also would like to thank the staff of the STAR program for providing this opportunity.

This material is based upon work supported by the S.D. Bechtel, Jr. Foundation and by the National Science Foundation under Grant No. 0952013 and Grant No. 0833353. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the S.D. Bechtel, Jr. Foundation or the National Science Foundation. This project has also been made possible with support of the National Marine Sanctuary Foundation.

The STAR program is administered by the Cal Poly Center for Excellence in Science and Mathematics Education (CESaME) on behalf of the California State University (CSU).

# References

- Abia, C. & Isern, J., 1997, MNRAS, 289, L11.
- Blanco et al., 1982, A&A, 115, 280.
- Böhm-Vitense, E., 1982, ApJ, 258, 628.
- Carpenter et al. 1997, ApJ, 486, 457.
- Dušek, J., Mikulášek, Z. & Papoušek, J., 2003, Contrib. Astron. Obs. Skalnaté Pleso, 33, 119.
- Evans, T.L., 2009, AIP Conf. Proc., 1094, 964.
- Groenewegen, M.A.T., van den Hoek, L.B., de Jong, T., 1995, A&A, 293, 381
- Judge, P.G. & Jordan, C., 1991 ApJ, 77, 75.
- Johnson, H.R. & Luttermoser, D.G., 1987, ApJ, 314, 329.
- Kenyon et. Al, 1991, ApJ, 101, 637.
- Kerschbaum, F. & Hron, J. 1994, A&A, 106, 397.
- Kerschbaum, F. & Oloffson, H. 1999, A&A, 138, 299.
- Knapp et al., 1999, A&A, 351, 97.
- Luttermoser, D.G. 2000, ApJ, 536, 923.
- Luttermoser, D.G. & Mahar, S. 1998, ASPC, 154, 1613.
- Morrissey et al., 2007, ApJ, 173, 682.
- Munari, U. & Buson, L.M., 1994, A&A, 287, 87.
- Nakashima, J., 2005, ApJ, draft.
- Nakashima, J., 2006, IAU, 234.
- Olivier, E.A. & Wood, P.R. 2003, ApJ, 584, 1035.
- Percy, J.R., Wilson, J.B., Hengy, G.W., 2001, ASP, 113, 983.
- Ramstedt et al., 2012, A&A, 543, A147.
- Riemers, D. & Cassatella, A. 1985, ApJ, 297, 275.
- Sanad et al., 2009, ApJ, 137:3479.
- Sahai, R. 2012, GALEX proposal #54.
- Sahai, R. et al., 2008, ApJ, 689, 1274.
- Shore, S.N. & Aufdenberg, J.P. 1993, ApJ, 416, 355.
- Stencel, R.E., 1982, 392, 137.
- Wallerstein et al., 2011, A&A, 535, A101
- Wood & Karovska, 2000, ApJ, 535, 304.