

Search for Electromagnetic Counterparts to LIGO-Virgo Candidates: Expanded Very Large Array[†] Observations

Joseph Lazio^{1†}, Katie Keating^{2¶}, F. A. Jenet^{3||}, and N. E. Kassim^{4††}

¹Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr.,
M/S 138-308, Pasadena, CA 91109, USA
email: Joseph.Lazio@jpl.nasa.gov

²National Research Council, Naval Research Laboratory, 4555 Overlook Avenue SW,
Washington, DC 20375-5351 USA
email: katie.mae.keating@gmail.com

³Center for Gravitational Wave Astronomy, University of Texas at Brownsville, Brownsville,
TX 78520, USA
email: merlyn@phys.utb.edu

⁴Remote Sensing Division, Naval Research Laboratory, 4555 Overlook Avenue SW,
Washington, DC 20375-5351 USA
email: Namir.Kassim@nrl.navy.mil

Abstract. This paper summarizes a search for radio wavelength counterparts to candidate gravitational wave events. The identification of an electromagnetic counterpart could provide a more complete understanding of a gravitational wave event, including such characteristics as the location and the nature of the progenitor. We used the Expanded Very Large Array (EVLA) to search six galaxies which were identified as potential hosts for two candidate gravitational wave events. We summarize our procedures and discuss preliminary results.

Keywords. Gravitational waves, methods: observational, radio continuum: general

1. Gravitational Wave Astronomy and the Time Domain

Gravitational waves (GWs) are fluctuations in the spacetime metric, equivalent to electromagnetic waves resulting from fluctuations in an electromagnetic field. Because of the weakness of the gravitational force, however, a laboratory demonstration of GWs comparable to Hertz's demonstration of electromagnetic waves is not possible. Indeed, the characteristic scale for the luminosity of a GW source is $L_0 = 2 \times 10^5 M_\odot c^2 \text{ s}^{-1}$, indicating immediately that the generation of GWs will occur in astrophysical environments in which large masses are moving at high velocities.

Precise timing of pulses from the radio pulsar PSR B1913+16, which is one member of a double neutron star system, has already revealed indirect evidence for GWs (Hulse & Taylor 1975; Weisberg et al. 2010). In this system, the rate of orbital period decay as

[†] The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

[‡] Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration.

[¶] A portion of this work was performed while K. K. held a National Research Council-NRL Research Associateship.

^{||} This work was supported by a CAREER grant from the National Science Foundation (award AST 0545837).

^{††} Basic research in radio astronomy at the NRL is supported by 6.1 Base funding.

predicted by general relativity is consistent with that observed from the pulsar timing measurements. Since the discovery of that system, other pulsars in neutron star-neutron star binaries have also been discovered, and the predicted level of orbital period decay from general relativity remains consistent with that from the measurements.

From the standpoint of time domain astronomy, many of the other predicted sources of GWs are rapidly time varying phenomena. These include the mergers of compact objects (neutron star-neutron star, neutron star-black hole, and black hole-black hole mergers), asymmetric supernovae, rapidly rotating asymmetric neutron stars, and exotic objects such as oscillating cosmic strings.

While evidence for GWs remains indirect, the promise of direct detection of GWs has excited considerable international interest. In Europe, the AstroNet *A Science Vision for European Astronomy* posed “Can we observe strong gravity in action?” as a key question for this decade while, in the U.S., gravitational wave astronomy was identified as a scientific frontier discovery area in the *New Worlds, New Horizons in Astronomy and Astrophysics* Decadal Survey. In this respect, GW astronomy is similar to the experience in opening up new spectral windows in the electromagnetic spectrum. As each new electromagnetic spectral window has been opened, there have entirely new classes of sources discovered. Indeed, one of the most surprising results of GW astronomy would be if *no* new classes of sources were discovered as this new window on the Universe is opened.

2. Electromagnetic Counterparts to Gravitational Wave Events

Supernovae have a well-known electromagnetic signature, and asymmetric supernovae are predicted to be gravitational wave emitters. It is therefore natural to anticipate that other gravitational wave emitters might also display electromagnetic counterparts. Moreover, the identification of electromagnetic counterparts to GW events would have a number of benefits, such as

- Precise localization of the event, potentially at the sub-arcsecond level, is possible via electromagnetic observations. Such localization may be crucial in understanding the nature of the event (e.g., in the nucleus of a galaxy vs. at its outskirts).
- The characteristics of the electromagnetic counterpart, such as its spectrum, will likely constrain the environment or progenitor or both of the GW event.

In many respects, determining the electromagnetic counterpart to a GW event is analogous to determining the (electromagnetic) spectrum of a transient discovered in one band and then followed up at others (e.g., a gamma-ray burst discovered at gamma-ray wavelengths, then followed up at X-ray, optical, and radio wavelengths).

Our focus here is on radio wavelength counterparts, motivated by several considerations:

- Non-thermal, high energy particles often produce radio wavelength emission easily, particularly in the presence of a magnetic field (e.g., cyclotron and synchrotron emission).
- Radio wavelengths can obtain precise astrometry, in the best cases obtaining positions at the milliarcsecond level.
- Radio wavelengths are unaffected by dust obscuration, either from the immediate environment of the event or from intervening objects.
- Radio telescopes can observe during the day, offering rapid followup.
- If a GW event also produces a radio burst or pulse, the propagation of this pulse will be delayed by its propagation through the ionized interstellar (and intergalactic) medium. Such delays can be minutes to hours, depending upon the electron column density along the line of sight, but they potentially allow for detailed followup of the burst.

3. LIGO-Virgo Observations

The radio wavelength observations that we describe below are based on coordinated observations between the Laser Interferometric Gravitational-wave Observatory (LIGO) and the Virgo that occurred during the Autumn of 2010 (LIGO Scientific Consortium & Virgo Collaboration 2011). LIGO has two elements, one located in Hanford, WA, USA, and the other in Livingston, LA, USA, while Virgo has one element located near Pisa, Italy. Together they form a 3-element interferometer.

The LIGO-Virgo interferometer measures time differences of arrival. Analysis of test waveforms injected into the LIGO-Virgo processing pipeline indicate that a candidate's position can be localized only to a region of order 10 deg^2 . Ordinarily, such a large uncertainty region could not usefully be searched for a radio counterpart with the current generation of telescopes, because their fields of view are too small. However, the most likely sources that the LIGO-Virgo interferometer could detect at reasonable signal-to-noise ratios would have occurred within 40 Mpc. If one assumes that a candidate event is associated with a galaxy, then the typical number of galaxies within an GW candidate certainty region is only three. This small number of galaxies can be usefully searched.

4. Expanded Very Large Array Observations

The Expanded Very Large Array (EVLA) is a 27-element radio interferometer operating between 1 and 50 GHz. It has been the focus of a recent major upgrade, which is nearly complete, and it is being commissioned with science programs now well established. For the purpose of these observations, the EVLA offers a number of attractive features.

- The wide frequency (wavelength) coverage potentially allows “tuning” of the observations to a frequency well matched to the expected physics. In this case, we observed at 5 GHz ($\lambda 6\text{cm}$), at which both expanding synchrotron fireballs and relativistic jets are likely to be detectable.

- At our observational frequency, the nominal field of view is approximately $7'$, well matched to the size of most local galaxies. In practice, the field of view is usually defined as the region over which the antenna response is at least half of its peak value; sources outside of the nominal field of view can still be detected, provided that they are sufficiently strong to compensate for the decreased antenna response. Accordingly, in order not to miss a potential candidate, we imaged a much larger region (typically $30'$).

- The angular resolution of the EVLA can be adjusted by moving the individual elements. During most of our observations, the obtained angular resolution was about $4''$, which provides about $0''.4$ localization (equivalent to 20 pc at 10 Mpc), for a reasonable signal-to-noise ratio.

We have now conducted three epochs of observations for each of the two LIGO-Virgo candidates, observing all of the nearby galaxies within the uncertainty region. Figure 1 shows the field around one of the galaxies. We typically detected six sources in the field of each galaxy. This small number of sources is consistent with the number of extragalactic sources expected (Windhorst 2003) but does not exclude the possibility that one of the radio sources is a counterpart to the LIGO-Virgo candidate.

Data acquisition and reduction of all three epochs for both candidates has only recently been concluded. In assessing the reality of any potential radio wavelength counterpart to either LIGO-Virgo candidate, other potential sources of variability must also be considered. On the time scales and cadence of our observations, it is unlikely that intrinsic variability of any active galactic nuclei (AGN) in the field of view would represent a source




Figure 1. Radio wavelength ($\lambda 6\text{cm}$) sources detected in the field of view of one of the galaxies representing a potential host for a candidate LIGO-Virgo gravitational wave event. Crosses represent the detected sources, and the circle represents the approximate “field of view,” defined as the half-power point of the antenna response.

of contamination. However, at these wavelengths, refractive interstellar scintillation due to the Galaxy’s interstellar medium is a potential source of contamination (i.e., an AGN unrelated to the LIGO-Virgo candidate might show variability on the time scales of our observations).

5. Future

Both LIGO and Virgo are currently undergoing upgrades. When they resume observation (~ 2014), they will be more sensitive and able to probe to larger distances. This increased GW sensitivity will require a change in observing strategy. A much larger number of galaxies could now be hosts, and a search of all of them with the EVLA would likely be quite time consuming (see also Metzger & Berger 2011). An approach similar to current followup of supernovae could, however, be profitable, namely, if an optical counterpart is found, then the EVLA could be used to assess the radio wavelength properties of the counterpart.

Later in the decade, a number of other radio wavelength facilities are likely to be available, which will present additional opportunities. Among these facilities are the Low Frequency Array (LOFAR), which could conduct wide-field of view “blind” searches at meter wavelengths (30–240 MHz) for northern hemisphere counterparts; the Karoo Array Telescope (MeerKAT), which could conduct southern hemisphere followup observations similar to those of the EVLA; and the Australian Square Kilometre Array Pathfinder (ASKAP), which could conduct wide-field of view “blind” searches at decimeter wavelengths (~ 1 GHz) for southern hemisphere counterparts.

References

- Hulse, R. A., & Taylor, J. H. 1975, *ApJ*, 195, L51
 LIGO Scientific Collaboration & Virgo Collaboration, 2011, arXiv:1109.3498
 Metzger, B. D., & Berger, E. 2011, *ApJ*, in press; arXiv:1108.6056
 Weisberg, J. M., Nice, D. J., & Taylor, J. H. 2010, *ApJ*, 722, 1030
 Windhorst, R. A. 2003, *New Astron. Rev.*, 47, 357