

Analysis of Multiscale Ultrahigh Resolution (MUR) Sea Surface Temperature Data Sets

Robin Sehler

CSU STAR 2011

Mentor: Dr. Jorge Vazquez


Co-mentor: Dr. Edward Armstrong

Project Outline

- Objective
- Background
- Methods
- Data
- Analysis
- Conclusion
- Acknowledgements


Project Objective

- To determine which spatial resolution of satellite-derived MUR version 2 data best reflects ocean buoy sea surface temperatures:
 - 1 km spatial resolution (i.e. HIGH RESOLUTION)?
 - 9 km spatial resolution (i.e. LOW RESOLUTION)?





Background: MUR Version 2

- MUR is one of NASA's newest SST products, developed by Dr. Mike Chin
- MUR combines data taken from both
 - Infrared radiation sensors (limited by cloud coverage)
 - Microwave radiation sensors (limited by resolution)



Background: GHRSSST

- The study is a part of an international effort by the Group for High Resolution Sea Surface Temperature (GHRSSST) to provide high resolution SST data in a uniform format and with a complete description of errors (e.g. bias, standard deviation)
- GHRSSST data includes both NASA/NOAA sensors which are used to create the MUR data:
 - Moderate Resolution Imaging Spectroradiometer (MODIS)-NASA
 - The Advanced Very High Resolution Radiometer (AVHRR)-NOAA
 - Advanced Microwave Scanning Radiometer - Earth Observing System(AMSRE)-NASA





Why is this the study of SST important?

- The National Oceanic and Atmospheric Administration (**NOAA**) has defined SST as an “essential climate variable” (ECV)
- In order to study climate change, scientists must have climate data that does not have a bias between buoy-derived SST and satellite-derived SST.




Methodology

Use three IDL validation codes to compare mean bias and standard deviations of low resolution and high resolution data over the period of April 1 – December 31, 2009, for the following oceanic regions

- Oceans surrounding North and Central America
- The California Coast (major region of upwelling)
- The Gulf Stream (major region of ocean heat transport)




North and Central America




North and Central America




NCAMERICA (Drifting Buoys)



NCAMERICA (Drifting Buoys)




The California Coast




The California Coast




California Coast (Fixed Buoys)





California Coast (Fixed Buoys)



The Gulf Stream



The Gulf Stream



Conclusion

- Over the entire NCAMERICA region, including the oceans off the coast of California and in the Gulf Stream, the higher resolution dataset has consistently smaller bias and standard deviation values
- Off the coast of California, the largest biases and standard deviation values occur during times of upwelling.

Future Work

- Compare the high resolution and low resolution data sets using the same exact grids
- Look more carefully at the seasonal differences between HiRes/LowRes datasets in terms of their biases and standard deviations
- Better understand what is causing the bias to be +/- in certain geographic locations

Acknowledgements

- Mentor: Dr. Jorge Vazquez
- Dr. Edward Armstrong
- Dr. Toshio Mike Chin
- Grant Wagner
- CSU STAR
- NASA Jet Propulsion Laboratory
- Dr. Gerry Simila
- Martin Mathews
- Noyce Scholars Program

Acknowledgements

“This material is based upon work supported by the S.D. Bechtel, Jr. Foundation and by the National Science Foundation under Grant No. 0952013 and Grant No. 0733758. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the S.D. Bechtel, Jr. Foundation or the National Science Foundation.”

The End