# ACCESS concept study for the direct imaging and spectroscopy of exoplanetary systems



John Trauger (Jet Propulsion Laboratory / Caltech)
and the ACCESS Science and Engineering Team

Pathways towards Habitable Planets Conference
CosmoCaixa -- Barcelona -- 16 September 2009

(c) 2009 California Institute of Technology. Government sponsorship acknowledged

# ACCESS Science and Engineering Team

*John Trauger, Karl Stapelfeldt, Wesley Traub, John Krist*
*Dwight Moody, Eugene Serabyn, Dimitri Mawet, Laurent Pueyo, Stuart Shaklan*
*Curt Henry, Peggy Park, Rob Gappinger, Paul Brugarolas, James Alexander*
*Virgil Mireles, Olivia Dawson*
*(Jet Propulsion Laboratory)*

*Olivier Guyon (University of Arizona & Subaru Telescope)*

*Jeremy Kasdin, Bob Vanderbei, David Spergel (Princeton University)*

*Ruslan Belikov (NASA/Ames)*

*Geoff Marcy (UC Berkeley)*

*Robert Brown (STScI)*

*Jean Schneider (Paris Observatory)*

*Bruce Woodgate (NASA/Goddard)*

*Gary Matthews, Rob Egerman, Perry Voyer, Phillip Vallone*
*Jason Elias, Yves Conturie (ITT)*

*Ron Polidan, Chuck Lillie, Connie Spittler, David Lee*
*Reem Hejal, Allen Bronowick, Nick Saldivar (NGAS)*

*Mark Ealey, Tom Price (NGC/Xinetics)*

# *In a nutshell -- ACCESS*

- *A space telescope for the astrophysics of exoplanetary systems:*
    - *Direct imaging*
    - *Spectroscopy*
- *With a coronagraph to suppress the glare of starlight*
- *ACCESS asks: what coronagraphic mission is possible with technologies available today?*

# *Science objectives*

## *With emphasis on exoplanetary systems, possibly dynamically full, that harbor exoplanets, planetesimals, dust structures.*

- *Direct coronagraphic imaging and low-resolution (R=20) spectroscopy of exoplanet systems in reflected starlight, to include:*
- *Census of nearby known RV planets in orbits beyond ~1AU*
- *Search for mature exoplanet systems beyond the RV survey limits, including giant planets, super-earths, and possibly a dozen earth-mass planets*
- *Observe dust structures as an indicator of unseen planets and planetesimals*
- *Dust structure in the circumstellar environment as a probe of the life cycle of planetary systems: from young stellar objects to proto-planetary nebulae*



# *ACCESS = Actively-corrected Coronagraph Concepts for Exoplanetary System Studies*

- *Coronagraphic imaging and spectroscopy of exoplanetary systems in reflected starlight at visible wavelengths (450-900 nm)*
- *ACCESS is one of four medium-class exoplanet concepts selected by NASA for Astrophysics Strategic Mission Concept Studies*
- *Study compares performance and readiness of four major coronagraph architectures*
- *Defines a conceptual space observatory platform as the "level playing field" for comparisons among coronagraph types*
- *Also uses laboratory validation on JPL's HCIT as another "level playing field" for coronagraph hardware readiness*
- *Identifies a genre of scientifically compelling exoplanet mission concepts built upon mature (TRL6+) subsystem technologies*
- *Evaluates science reach of a medium-class coronagraph mission*

# *ACCESS observatory*

- *Observatory architecture is representative of the "best" available for exoplanet coronagraphy within the scope (cost, risk, schedule) of a NASA medium-class mission*
  - *visible wavelengths (500-900 nm) for smaller $\lambda/D$ and better IWA*
  - *single spacecraft (external occulter + telescope exceeds medium cost)*
- *In particular, all coronagraphs require an observatory system with*
  - *exceptional pointing control*
  - *exceptional wavefront stability and*
  - *active (deformable mirrors) wavefront control*
- *ACCESS requires systems with high technology readiness (TRL6+) for*
  - *reliable estimates of science capabilities and*
  - *reliable determinations of cost and schedule*
- *Baseline observatory architecture defines a capable platform for meaningful comparisons among coronagraph types.*

*ACCESS observatory:*
*1.5 meter - unobscured off-axis gregorian telescope*





# *ACCESS Discovery Space is critically dependent on observatory pointing control and wavefront stability*




*Pointing control and low-order wavefront stability*

*Wavefront stability*

# *ACCESS gamut of coronagraph types*







*The four major coronagraph types perform starlight rejection with combinations of phase and amplitude elements placed in focal and pupil planes. Best demonstrated laboratory contrast to date (March 2009) are indicated at right, while noting that significant improvements are expected this year as an outcome of active laboratory developments with well-understood technologies.*

# *ACCESS gamut of coronagraph types*







*The four major coronagraph types perform starlight rejection with combinations of phase and amplitude elements placed in focal and pupil planes. Best demonstrated laboratory contrast to date (March 2009) are indicated at right, while noting that significant improvements are expected this year as an outcome of active laboratory developments with well-understood technologies.*

# *Laboratory coronagraph contrast and stability demonstrate capability to detect exoplanets*






*Comparison of azimuthally averaged PSFs of (a) the star, with focal plane mask offset and Lyot stop in place; (b) the coronagraph field with all DM actuators set to equal voltages; (c) the coronagraph with DM set for a dark half-field; and (d) the result of simulated roll deconvolution with the set of 480 consecutive coronagraph images. PSFs of a nominal Earth and Jupiter are also indicated.* (Trauger & Traub, *Nature*, 12 April 2007, p771)

# ACCESS compares four major coronagraph types

## Lyot coronagraph




## Vortex coronagraph




## Shaped pupil coronagraph




*(Note: highlighted elements, including FSM, DMs, and pointing control system, are common to ALL four coronagraph types)*

## Pupil mapping coronagraph



# *Precision deformable mirrors provide wavefront control*








*Evolution of monolithic PMN deformable mirrors: left to right: 32x32 array, used for all HCIT milestones to date; 64x64 array to be installed on HCIT spring 2009; 48x48 array (also shown on JPL shake table) will be used to demonstrate TRL6 flight-readiness this year.*

*Mirror facesheets are fused silica, with surfaces polished nominally to* $\lambda/100$ *rms. Surface figure (open loop) is settable and stable to 0.01 nm rms over periods of 6 hours or more in a vacuum testbed environment. All DMs were delivered to JPL by Xinetics.*

# *Exoplanet completeness space*







*Examples of exoplanet search space. At left, a plot of integration times needed to detect Jupiter twins within 45 degrees of elongation from their parent stars, to S/N = 10, using the ACCESS Lyot coronagraph with an IWA = 2 lambda/D. At right, the number of planets detectable to S/N = 10, in integration times of one day or less, using the ACCESS Lyot coronagraph with an IWA = 2.5 lambda/D.*

# Science completeness metric: 2.0, 2.5, and 3.0 λ/D

Table 1: Number of nearby stars that can be surveyed for 5.2 AU Jupiters, IWA 3 l/D

| Coronagraph Type | Planet 45° from max elong | Planet 15° from max elong |
|---|---|---|
| Lyot | 117 | 175 |
| PIAA | 166 | 278 |
| Vortex | 135 | 204 |

Table 2: Number of nearby stars that can be surveyed for 5.2 AU Jupiters, IWA 2.5 l/D

| Coronagraph Type | Planet 45° from max elong | Planet 15° from max elong |
|---|---|---|
| Lyot | 153 | 218 |
| PIAA | 178 | 267 |
| Vortex | 154 | 228 |

Table 3: Number of nearby stars that can be surveyed for 5.2 AU Jupiters, IWA 2.0 l/D

| Coronagraph Type | Planet 45° from max elong | Planet 15° from max elong |
|---|---|---|
| Lyot | 170 | 230 |
| Vortex | 164 | 241 |

Note: Accurate PIAA wavefront control solution not available for this IWA

*Tabulation of the number of nearby stars that could be searched with various ACCESS coronagraphs to the depth of 10-sigma detections of Jupiter twins in each of six visits to the star over a period of 2.5 years. The green-highlighted row is the demonstrated coronagraph state of the art, 3.0 lambda/D with the Lyot coronagraph. The blue-highlighted rows represent coronagraph demonstrations that can realistically be achieved this year. Columns for 45 degrees from maximum elongation corresponds to an observational completeness of 50% or more in each visit, approaching 100% after six epochs spread over several years, and for systems with more than one major planet.*

# *All four coronagraph types have been tested in JPL's High Contrast Imaging Testbed*



# Lyot coronagraph demonstrations on the HCIT




THICKNESS-PROFILED NICKEL MASK

Nickel mask has been vacuum-deposited on a fused silica substrate. Attenuation profile was built up in a number of passes with a computer-controlled moving slit. The same mechanism will be used to superimpose a dielectric phase layer in future work.




Comparison of the prescribed transmittance profile with the measured profile of the mask pictured at left. Desired profile is the red curve, the measured profile is the blue curve.

*Recent contrast demonstrations in the HCIT:*
*IWA = 3 λ/D, 10% bandwidth, C = 1.2 e-9*
*IWA = 3 λ/D, 20% BW, C = 2.7 e-9*
*IWA = 4 λ/D, 10% BW, C = 6 e-10,*
*with 4th-order metallic or metal+dielectric 4th-order Lyot masks. All masks manufactured at JPL. Narrower (2.5 λ/D) Lyot masks, and circular masks will be manufactured this year.*

# Vector vortex coronagraph mask for HCIT experiments

*Recent contrast demonstrations in the HCIT:*
*IWA = 3 λ/D, 2% bandwidth, C = 2.0 e-7*
*with the first-ever charge-4 liquid crystal polymer vortex mask from JDSU (seen at left through crossed polarizers). Close agreement between HCIT performance and models predict that reduction of internal reflections and multilayer achromatization will lead to contrast ~1e-9 with a 20% bandwidth, to be attempted by the end of this year.*



# HCIT laboratory setup for pupil mapping demonstrations

*To date, the best contrast result for a pupil mapping (PIAA) coronagraph: IWA = 1.6 λ/D, monochromatic, C = 2.2e-7 was achieved by Guyon (2009) at the Subaru Observatory laboratories. A new PIAA system, commissioned by NASA/Ames and manufactured by Tinsley, has completed a preliminary experiments in vacuum.*




# Shaped pupil coronagraph experiments with HCIT



*At left, the transmittance profile of a representative shaped pupil apodization (black indicates opaque). This "Ripple 3" design achieved: IWA = 4 λ/D, 10% bandwidth, C = 2.4 e-9 on the HCIT (Belikov et al. 2007). Smaller inner working angles are possible with the use of a pair of DMs*

# *ACCESS gamut of coronagraph types*







*The four major coronagraph types perform starlight rejection with combinations of phase and amplitude elements placed in focal and pupil planes. Best demonstrated laboratory contrast to date (March 2009) are indicated at right, while noting that significant improvements are expected this year as an outcome of active laboratory developments with well-understood technologies.*

# ACCESS




*ACCESS cost and schedule conform to guidelines for a NASA medium-class mission. The science mission would begin approximately 5 years after selection.*

# *Summary*

- *The ACCESS study considered the relative merits and readiness of four major coronagraph types, and hybrid combinations.*
- *Using demonstrated high-TRL technologies, the ACCESS minimum science program surveys the nearest 120+ AFGK stars for exoplanet systems, and surveys the majority of those for exozodiacal dust to the level of 1 zodi at 3 AU.*
- *Follow-up with R=20 spectrophotometry.*
- *Technology demonstrations in the coming year are expected to further enhance the science reach of an ACCESS mission, in advance of a NASA AO for a medium class mission.*
- *The study also identifies areas of technology development that would advance the readiness of the major coronagraph types in the coming 5 years.*

# End