Comparing spacecraft and laboratory analog spectra to determine the compositions of Phobos and Deimos

Abigail Fraeman, JPL

UCLA Asteroid-Meteorite Connection Workshop

April. 21-22, 2016

Image credit: NASA/JPL/University of Arizona

Phobos and Deimos Quick Facts



	Phobos	Deimos
Size	26.1 x 22.8 x 18.3 km	10.4 x 12.2 x 15 km
Density	1.86 g/cm^3	1.49 g/cm^3
Orbital period	7.66 hr	30.35 hr
Semi-major axis	9,377 km	23,460 km
Eccentricity	0.0151	0.00033
Inclination	1.093°	0.93°

Murchie et al., 2015 and ref. therein



Image credit: NASA/JPL/University of Arizona

Origin Questions

EXTERNAL

Captured asteroids
Burns, 1978

IN SITU

Formation by co-accretion
Safronov et al., 1986



Giant Impact Hypothesis
Craddock et al., 2010

Expected compositions

- Captured primitive bodies: compositions typical of material found in primitive meteorites
- Form by late stage co-accretion with Mars: compositions consistent with bulk Mars chondritic mafic mineralogy, e.g. ordinary chondrites
- Form from differentiated Mars (impact hypothesis): compositions with mafic minerals similar to basaltic Martian crust

Murchison Carbonaceous Chondrite

Image Credit: Randy Korotev

Unnamed Ordinary Chondrite

Image Credit: Randy Korotev

Adirondack

Image Credit: NASA/JPL/Cornell

Expected Composition

Table 1
Compositions predicted by different models for the origin of Phobos.

Origin Hypothesis	Composition predicted	Elemental abundances	Mineral abundances
Capture of organic- and water-rich outer solar system body	Ultra-primitive composition; Tagish Lake is the best known analog (Brown et al., 2000)	High C; high Zn/Mn; high S; composition possibly unique from known meteorites	Abundant phyllosilicates; carbonates and organic phases; anhydrous silicate phases rare
Capture of organic- and water-poor outer solar system body	Anhydrous silicates plus elemental carbon (Emery and Brown, 2004)	High C; Mg/Fe ratio ~2–4; bulk composition unlike any meteorite analogs	Anhydrous, med. Fe (20–40%) pyroxene+olivine; abundant amorphous carbon or graphite?
Capture of inner solar system body	Composition like common meteorites (e.g. ordinary chondrites) (Brearley and Jones, 1998)	Mg/Si ~0.8–1, Al/Si ~0.05–0.1; Zn/Mn & Al/Mn ratios separate known meteorites; likely low C	Low carbonates, phyllosilicates; pyroxene, olivine probably in range of known meteorites
Co-accretion with Mars	Bulk Mars; similar to ordinary chondrites but specific SNC-derived comp. (Wanke and Dreibus, 1988)	Mg/Si, Al/Si, Fe/Si indicative of bulk Mars; low C; Zn/Mn, Al/Mn like ordinary chondrites	Anhydrous silicates with Fe, Mg of bulk Mars; low abundance of C-bearing phases
Giant impact on Mars	Evolved martian crust or mantle, like SNC meteorites, Mars rocks or soil (McSween et al., 2009)+impactor	High Al/Si, Ca/Si, lower Fe/Si, Mg/Si indicative of evolved igneous materials	Evolved, basaltic mineralogy consistent with many datasets for Mars

Murchie et al., 2014

Composition: View in Mid-2000s

- Spectral measurements showed Phobos had "blue" and "redder" material; Deimos looks like Phobos
- Redder material looked like D-type primitive asteroids
- No clear diagnostic evidence for mineral absorptions



Thomson et al., 2011



Rivkin et al., 2002

Spectrometers Orbiting Mars: OMEGA and CRISM

	OMEGA	CRISM
Spacecraft	Mars Express	Mars Reconnaissance Orbiter
Wavelength Range	0.35 – 5.0 μm	0.4 – 3.9 μm
Spectral Resolution	0.007 to 0.02 μm/channel	0.00655 μm/channel
Pixel Angular Size	1.2 mrad	0.0615 mrad

OMEGA



Bibring et al., 2004

CRISM



Murchie et al., 2007

Mars Express (and MRO) Observations



Image credit: ESA/DLR

OMEGA Orbital Datasets

OMEGA: High Spectral Resolution



2200 m/pix 200 m/pix 1170 m/pix 800 m/pix 120 m/pix 380 m/pix

HRSC: High Spatial Resolution Context

75 m/pix 7 m/pix ~40 m/pix ~27 m/pix 3.7 m/pix 9 m/pix

CRISM Orbital Datasets

CRISM: High Spectral Resolution

Phobos

350 m/pixel

Deimos



1200 m/pixel

HiRISE: High Spatial Resolution Context



6.8 m/pixel

20 m/pixel

Summary Spectra

OMEGA Phobos



CRISM Phobos



CRISM Deimos





Different Lighting Conditions



CRISM Phobos | OMEGA Phobos | CRISM Deimos



Incidence Angle (i)

Emergence angle (e)

Phase angle (g)



Modeling Effects of Viewing Geometry on Reflected Solar Radiance

Hapke's Model for Radiance Factor

$$r(i,e,\alpha)=\frac{w}{4\pi}\frac{\mu_0}{\mu_0+\mu}\left[(1+B(g))p(g_1)+H(w,\mu_0)H(w,\mu)-1\right]\cdot S(i,e,g,\theta)$$

Single scattering alebdo: ratio of scattering to scattering absorption efficiencies for a single event

Related to viewing geometry

Approx. for scattering between particles

Related to surface roughness

Measured/Calculated: r, i, e, g, μ_0, μ
Unknown: w, g_1, Θ

Fraeman et al., 2012

Range of OMEGA Lighting Geometries





Fraeman et al., 2012

Solve for Single Scattering Alebdo at Every Pixel





CRISM Phobos

OMEGA Phobos

CRISM Deimos

Fraeman et al., 2012

Recast to Any Lighting



Fraeman et al., 2012

Recast to Any Lighting



Fraeman et al., 2012

Space Weathering of Carbonaceous Meteorites

- Telescopic studies of asteroids of different ages suggest opposite trends in spectral slope with age (Nesvorny, et al., 2005; Lazzarin, et al., 2006; Kaluna et al., 2015)
- Experiments on carbonaceous chondrites suggest spectral properties might darken and redden like lunar samples, although details vary between samples (Gillis-Davis et al., 2015; Matsuoka et al., 2016)
- Haven't (yet) examined returned sample from carbonaceous asteroid



Fig. 1 Reflectance values for Allende, Murchison and Graphite as a function of space weathering.

Gillis-Davis, J. et al., LPSC 2015

Searching for Features in CRISM Data



Fraeman et al., 2014

Broad Feature at 0.65 µm

Phobos Deimos





Fraeman et al., 2014

OSIRIS Data from Rosetta

- OSIRIS on Rosetta spacecraft also collected spectra from Phobos during Mars flyby
- Data were consistent with previous studies



Figure 3. Scaled OSIRIS WAC and NAC data together with the best fitting Models 1 and 2.

Pajola et al., 2013

Jan. 2016 OMEGA Observation
53 km above Phobos (ORB 15260)

Red Unit



Blue Unit







Spectra courtesy B. Gondet

Feature at 2.8 μm





Fraeman et al., 2014

Similarity to Low-Albedo Asteroids

0.65 μm Feature



2.8 μm Feature



Vilas et al., 1998

Rivkin et al., 1996; Rivkin et al., in prep

Scenario 1: Phyllosilicate





Fraeman et al., 2014

Scenario 2: Space Weathering





Fraeman et al., 2014

Thermal Infrared Measurements



Figure 1. Topographic map of Phobos with the approximate footprints of several TES spectra of Phobos from orbit 551. ICK = incremental counter keeper. DET = detector.

Surface hydration?

Indicates low Si content if space weathered

Carbonates + silicates



Figure 3. Average TES Phobos spectrum, compiled from the 88 high quality spectra shown in Figure 2.

Glotch et al., 2015

Key Conclusions

- Phobos and Deimos are darker than even most space weathered materials
- Likely have carbonaceous chondrite-like compositions because they lack mafic absorptions and are spectrally similar to CM carbonaceous chondrites or Tagish Lake
- Pair of spectral features observed similar to those on low albedo asteroids
- To argue that moons formed *in situ* rather than by capture of primitive bodies requires carbonaceous materials to have been added to the Martian system during accretion or a late stage impact

Phobos Surface Geology



Wählisch, M., et al., (2013), http://dx.doi.org/10.1016/j.pss.2013.05.012i

Want to learn more?

- http://sservi.nasa.gov/event/planetary-evolution-phobos-and-deimos/

- The "Science and Exploration of Phobos and Deimos" series was jointly organized and led by SSERVI teams at University of Central FL and Brown University/MIT with many SSERVI-affiliated institutions participating.
- 13 recorded seminars given by leaders in the field with associated discussion.
- Also includes great reading list.

Space Weathering

- Catch all phrases for processes that affect spectral properties of materials due to their exposure in the space environment
- Biggest drivers are solar-wind irradiation and micrometeorite impacts
- Effects of space weathering known to be different for different solar system bodies
  - Moon = production of nanophase Fe that darkens and reddens, observed in returned lunar samples (Hapke, 2001; Pieters, et al., 2000)
  - Itokawa = production of FeS as well (Noguchi et al., 2011)
  - Vesta = little evidence for space weathering (Pieters et al., 2012)

Composition to help Constrain Internal Structure

- Once we know composition, we can use it to make assumptions about other physical properties, like internal structure
- Helps constrain knowledge of dissipative orbital properties, which is key to understanding past evolution of orbits (e.g. Lambeck 1979; Mignard 1981)
- See Rosenblatt, 2011 for thorough review of links between internal structure and origin

Internal Structure



DTM of Phobos from Willner, et al. 2013 created using stereo-photogrammetric analyses of 100m/pixel HRSC and Viking Orbiter data

- MRO and MEx allow for improved estimates for Phobos mass and volume (Andert, et al., 2011; Jacobson, et al., 2010, Willner, et al., 2013)
- Ph density: 1.86 ± 0.013 g cm^{-3}
- D density: 1.490 ± 0.190 g cm^{-3}



Rosenblatt, 2011

Internal characteristics & surface manifestations



Fig. 1. Viking Orbiter images showing (a) Phobos' well-organized pattern of grooves [20] and (b) Deimos' surface dominated by bright streaks due to mass wasting, with a large depression in the south polar region that is thought to be a large impact crater [21].

Body	R_m, km	Assumed composition	Est. macro-porosity	Est. total porosity	Coherent ridge/ groove patterns?	Craters >1.3 R_m?	Fragment-dominated surface?	Refs.
Coherent to fractured but coherent								
Lutetia	49	E	≤15%	≤17%	Y	N	N	25,26
Phobos	11.1	CM	17±4%	35±3%	Y	N	N	13,17,18
Eros	7.3	LL	17±7%	24±4%	Y	N	N	7,10
Fractured but coherent to transitional								
Mathilde	26.5	CI	18±14%	55±9%	insufficient resolution	Y	insufficient resolution	5,22
Ida	15.7	LL	20±20%	26±16%	Y (only regionally)	N	N	27,28
Loosely consolidated								
Deimos	6.2	CM	34±11%	48±8%	N	Y	Y	14,16-18
Itokawa	0.18	LL	41±8%	46±6%	N	N	Y	8,9,11

Table 1. Sizes, assumed compositions, estimated porosities calculated using meteorite densities reported by [4], and morphology of asteroids <110 km in diameter having well-determined densities and coverage by spacecraft images. R_m=mean radius.

Murchie et al., 2013

Future Exploration



Image Credit: Victor Dang and the Caltech Space Challenge Explorer Team

Spacecraft Observations

- Mars flybys
  - Mariners 4, 5, and 6
  - Rosetta
- Mars orbiters
  - Mariner 9
  - Viking orbiters 1 & 2
  - Phobos 88
  - Mars Global Surveyor
  - Mars Odyssey
  - Mars Express (MEx)
  - Mars Reconnaissance Orbiter (MRO)
- Surface of Mars
  - Viking Landers 1 & 2
  - Pathfinder
  - Spirit & Opportunity
  - Curiosity

Phobos and Deimos as seen from the Martian surface by Curiosity on August 1st, 2013.

Image credit: NASA/JPL-Caltech/Malin Space Science Systems/Texas A&M Univ.

List source: Duxbury et al., 2014

Red and Blue Units





Fig. 17. Schematic illustrative diagram of a profile from the eastern rim of the crater Stickney towards the crater center and then to its northern rim. Rim structure and bedrock deformation has been omitted. The stages of formation of the observed morphology and distribution of the red and blue units are shown: (a) stage 1, (b) stage 2 and (c) stage 3 for two cases (see text). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Basilevsky et al., 2014