



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# NEXUS

# Big Data Analytics

**Thomas Huang**

NASA Physical Oceanography Distributed Active Archive Center (PO.DAAC)

Jet Propulsion Laboratory  
California Institute of Technology  
4800 Oak Grove Drive  
Pasadena, CA 91109-8099  
United States of America

© 2016 California Institute of Technology. Government sponsorship acknowledged.  
Reference herein to any specific commercial product, process, or service by trade name,  
trademark, manufacturer, or otherwise, does not constitute or imply its endorsements by the  
United States Government or the Jet Propulsion Laboratory, California Institute of Technology.

THUANG/JPL

2016 ESIP Federation Summer Meeting



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

“Traditional data management products are built on top of a scale-up architecture that cannot handle petabyte-scale applications...”

– *Why Traditional Data Management Fails in the Eras of NoSQL and Big Data*

“It’s not enough to just collect and store a lot of information; it’s what you do with the insights and learnings from that data that have real, measurable impacts on a business.”

– *The Art of Data Management*

“The Big Data technologies enable a whole new class of insight discovery, against data we’ve either not had access to before or not had the tools to efficiently process such as high volume time-series analysis or cluster analysis on a traditional RDBMS.”

– *Of Course Big Data Will Change the “Traditional” Data Warehouse*

“Climate change research is now ‘big science’, comparable in its magnitude, complexity and societal importance to human genomics and bioinformatics.”

– *Database bonanza*





National Aeronautics and  
Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# but ... How Do We Get There?




There





# Traditional Data Analysis



- Depending on the data volume (size and number of files)
- It could take many hours of download – (e.g. 10yr of observational data could yield thousands of files)
- It could take many hours of computation
- It requires expensive local computing resource (CPU + RAM + Storage)
- After result is produced, purge downloaded files

## Observation

- Traditional methods for data analysis (time-series, distribution, climatology generation) yield poor performance
- Performance suffers when involve large files and/or large collection of files
- A high-performance data analysis solution must be free from file I/O bottleneck



# NEXUS Deep Data Analytics: One-Minute Summary

**NEXUS** is an emerging technology developed at JPL

- A Cloud-based/Cluster-based data platform that performs scalable handling of observational parameters analysis designed to scale horizontally by
- Leveraging high-performance indexed, temporal, and geospatial search solution
- Breaks data products into small chunks and stores them in a Cloud-based data store

## Data Volumes Exploding


- SWOT mission is coming
- File I/O is slow

## Scalable Store & Compute is Available

- NoSQL cluster databases
- Parallel compute, in-memory map-reduce
- Bring Compute to Highly-Accessible Data (using Hybrid Cloud)

## Pre-Chunk and Summarize Key Variables

- Easy statistics instantly (milliseconds)
- Harder statistics on-demand (in seconds)
- Visualize original data (layers) on a map quickly




## Built with open source technologies

- Apache Solr
- Apache Cassandra
- Apache Spark/PySpark
- Apache Mesos/YARN
- Apache Kafka
- Apache Zookeeper
- Tornado
- Spring XD
- EDGE




# Analytics & Summarization Stack Using Two Scalable Database Technologies





# Deep Data Computing Environment (DDCE)

- ETL System – Ingest and stage data
- Deep Data Processors – metadata, statistics, and tiles
- Index and Data Catalog – horizontal-scale geospatial search and tile retrieval
- Analytic Platform – Spark-based domain-specific analytics
- Data Access – tile and collection-based data access
- Cloud Platform – portal and custom VMs



Credit JPL: T. Huang, B. Wilson, G. Chang, E. Armstrong, T. Chin


AIST-14: OceanXtremes





# The Notebook

## Interact with NEXUS using Jupyter Notebook



- **/capabilities**: list of capabilities
- **/chunks**: list data chunks by location, time, and datasets
- **/correlationMap**: Correlation Map
- **/datainbounds**: Matchup operation to fetch values from dataset within geographic bounds
- **/datapoint**: Matchup operation to fetch value at lat/lon point
- **/dailydifferenceaverage**: Daily difference average
- **/latitudeTimeHofMoeller**: Latitude Time Hovmoeller
- **/list**: list available datasets
- **/longitudeLatitudeMap**: Longitude Latitude Map
- **/longitudeTimeHofMoeller**: Longitude Time Hovmoeller
- **/stats**: Statistics (standard deviation, count, min/max, time, mean)




# Apache Spark and Cassandra

- **Apache Spark**

- In-Memory Map-Reduce framework
- Datasets partitioned across a compute cluster by key
- Resilient Distributed Dataset (RDD)
  - Fault-tolerant, parallel data structures
  - Intermediate results persisted in memory
  - User controls the partitioning to optimize data placement
- Rich set of operators on RDD's: *Map*, *Filter*, *GroupByKey*, *ReduceByKey*, etc.
- Computation is implicit (Lazy) until answers needed
- Uses YARN/Mesos


- **Apache Cassandra**

- Horizontal-scale NoSQL database
- Constant-time writes regardless of the size of data set grows
- No-single-point of failure architecture






# NASA Sea Level Change Portal – <https://sealevel.nasa.gov>




Data Comparison




Time-series




Latitude/Time Hovmoller



Data Analysis Tool




Longitude/Time Hovmoller





# On-The-Fly Analysis for Sea Level Rise Research




- Time Series
- Data Comparison
- Latitude/Time Hofmoeller
- Latitude/Longitude Time-Average



# AIST-14: OceanXtremes – Data-Intensive Anomaly Detection System

- Anomaly detection is a process of identifying items, events or observations outside the “norm” or expected patterns
- Current and future oceanographic missions and our research communities present us with challenges to rapidly identify features and anomalies in increasingly complex and voluminous observations
- Typically this is a two-stage procedure
  1. Determine a long-term/periodic mean (“climatology”)
  2. Deviations from the mean are searched. Step 1 could be omitted in cases where a climatology data set already exists.




## OceanXtremes System Architecture

Credit: T. Huang, E. Armstrong, G. Chang, T. Chin, B. Wilson




National Aeronautics and Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Xtremes Explorer



High Resolution Data Visualization for the Web




Data Analysis Workbench




National Aeronautics and  
Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California


# Daily Anomaly




Aug 02, 2012



Aug 02, 2013



Aug 02, 2014




Aug 02, 2015



National Aeronautics and  
Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# AIST-14: Distributed Oceanographic Matchup Service (DOMS)


- A distributed data service (a.k.a DOMS) to match satellite and in situ marine observations to support platform comparisons, cross-calibration, validation, and quality control
- Use Cases
  - Satellite Cal/Val and algorithm development
  - Decision support
    - Planning field campaigns
    - Real-time operational activities
  - Scientific investigation
    - Process studies
    - Model assimilation services
    - User friendly interface to support student research
  - Alternate matching
    - Satellite to satellite
    - Satellite/in situ to model





# Open Search for In-Situ Data

- **Common software interface for data search and retrieval (OpenSearch JSON)**
- **SAMOS @FSU**
  - Sample query: `curl "http://doms.coaps.fsu.edu/edge/samos?startTime=2012-08-01T00:00:00Z&endTime=2013-10-31T23:59:59Z&bbox=-45,15,-30,30"`
  - Integration with THREDDS and physical file archive
  - Full indexing of all relevant parameters: 54,911,530 documents (each with sst, wind, sss measurements)
- **ICOADS @NCAR**
  - Sample query: `curl "http://rda-db-icoads.ucar.edu:8890/ws/search/icoads?startTime=2012-08-01T00:00:00Z&endTime=2013-10-31T23:59:59Z&bbox=-45,15,-30,30"`
  - Integration with IVAD, MySQL server, hosts all ICOADS data exported from the original IMMA files
  - Full indexing of all relevant parameters: 517,140,830 documents (each with sst, wind, sss measurements)
- **SPURS Data Service**
  - Sample query: `curl "https://deepdata.jpl.nasa.gov/ws/search/spurs?startTime=2012-08-01T00:00:00Z&endTime=2013-10-31T23:59:59Z&bbox=-45,15,-30,30"`
  - 13,817,392 documents (each with sst, wind, and sss measurements)



Open Source, available on [Github.com](https://github.com/dataplumber/edge)  
<https://github.com/dataplumber/edge>





# NEXUS Performance Challenge

## **The Challenge: Show that NEXUS performs 2X or greater speed improvement compare to Giovanni – Sponsored by NASA/ESDIS**

- Dataset: TRMM Daily Precipitation (TRMM 3B42 Daily V7), 18 years, 6574 granule files, ~26GB
- Giovanni – backed by the popular NCO library
  - Highly optimized C/C++ library
- NEXUS
  - Apache Solr for spatial searches, metadata, and pre-computed statistics
  - Apache Cassandra for clustered data storage where granule data is partitioned into tiles
  - Apache Spark for data analytic platform


**Question:** With smaller tiles enables more parallelism. Does more parallelism yield faster performance?

- Divide daily granule into 16 tiles – 16 Executors and 16 Data Partitions
- Also experimented with more, smaller tiles - 545 tiles – 64 Executors and 128 Data Partitions



# NEXUS Performance Challenge Area Averaged Time Series


**Area Averaged Time Series:** The standard time-series plot produced by computing spatial averages over the user-selected area of a given variable for each time step within the user's range. Fill values do not contribute to the spatial averages. Each average value is then plotted against time to create the time-series output.



Global 18-year Area Averaged Map - Performance

## Observation


- On-the-fly data subsetting
- Giovanni took over an hour
- NEXUS, with smaller tiles, took less than 5min
- More tiles and more executors yield FASTER performance
- With 16-way parallel – NEXUS performs ~8X faster
- **With 64-way parallel – NEXUS performs ~15X faster**





# NEXUS Performance Challenge Global 18-Year Time Averaged Map


**Time Averaged Map:** The time averaged map shows data values for each grid cell within the user-specified area, averaged (linearly) over the user-specified time range as a map layer. Fill values do not contribute to the time averages.



Global 18-year Time Averaged Map - Performance

## Observation


- Global coverage
- Giovanni took about 3min
- NEXUS, with bigger tiles, took about 1min
- More tiles and more executors yield SLOWER performance
- With 16-way parallel – NEXUS performs ~2.8X faster





# NEXUS Performance Challenge Global 14-Year Correlation Map

**Global Correlation Map:** The correlation map calculates correlation coefficient using simple linear regression between two variables over time within each grid cell, producing two maps: one showing the correlation coefficient (R) and the other displaying the number of contributing (matching) samples in each grid cell. Any grid cell that contains fewer than three matched pairs over time will be assigned a fill value.




Global 14-year Correlation Map - Performance

## Observation

- Global coverage
- Giovanni took over 10min
- NEXUS, with bigger tiles, took a little over 1min
- More tiles and more executors yield SLOWER performance
- With 16-way parallel – NEXUS performs ~7X faster

Correlation of Precipitation Rate daily 0.25 deg. [TRMM TRMM 3B42 daily v7] mm/day vs. Ne...



Correlation of Precipitation Rate daily 0.25 deg. [TRMM TRMM 3B42 daily v7] mm/day vs. Near-Real-T...  
 0.0 0.2 0.4 0.6 0.8 1.0  
 Data Min = 0.0, Max = 1.0, Mean = 0.9



# Key Takeaways

- Global analysis - NEXUS out performs Giovanni with larger tiles and limit parallelism
- Area analysis - NEXUS out performs Giovanni with finer tiling scheme
- Another observation: NEXUS is a fast data subsetter
- More parallelism  $\neq$  Faster performance
  - For global analytics, more smaller tiles actually yields slower performance
  - Why? Scheduling, data transport, data queries, etc.
- Big Data  $\neq$  Cloud Computing
- Big Data is not a new computing problem. Cloud Computing opens up new approaches in tackling Big Data
- While Cloud Computing has many benefits, it only plays a part in the overall Big Data architecture
  - Process, Information Model, Technologies, etc.
- Apply Cloud Computing where it make senses
  - Data-Intensive Science, Cost reduction, Service reliability, etc.
- Part of the architectural design involves modernizing existing software solutions in order to
  - Truly leverage the elasticity of the Cloud
- It makes sense to bring the computing close to the data - On-premise Cloud (currently)
  - Need local experts
  - Governance
- Use automation deployment – Puppet, Chef, Salt
- Many technologies are mature in their standalone context
  - It doesn't mean they are high TRL when integrated into our domain-specific architecture
- Look into Open Source Solutions before build your own
- **NEXUS Current Activities**
  - It is still actively being developed and more to come
  - Part of the AIST Data Container Study



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# July 14, 2016

\*\*\*\* APPROVED \*\*\*\*

The software known as NEXUS: Deep Data Platform (NTR-50157) has been approved for release as open source. You are authorized to upload the software to open source repository when you are ready to do so.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# IN008: Big Data Analytics



[https://agu.confex.com/agu/fm16/preliminaryview.cgi/  
Session13144](https://agu.confex.com/agu/fm16/preliminaryview.cgi/Session13144)

## IN008: Big Data Analytics

[Submit an Abstract to this Session](#)

**Session ID#:** 13144

**Session Description:**

Big Data pose great challenges for Earth and Space sciences. Cloud Computing emerged as a promising solution for supporting Big Data analytics in areas such as climate science, ocean science, atmospheric science, planetary science, and other geoscience domains for model simulation, data management, information mining, decision support, and visualization. This session is to capture the latest on applying Cloud Computing for Big Data Analytical problems in all Earth and space domains. Topics include demonstration, studies, methods, solutions and solution discussion on  
Solutions for big data analytics  
Big data mining  
Application of open source technologies  
Automated techniques for data analysis  
Browser-based data analytics and visualization  
Real time decision support  
Contributions that fuse participatory social learning into the Geoscience R&D processes are also welcome.

**Primary Convener:**

**Thomas Huang**, NASA Jet Propulsion Laboratory, Pasadena, CA, United States

**Conveners:**

**Chaowei Phil Yang**, George Mason University Fairfax, Fairfax, VA, United States, **Tiffany C Vance**, NOAA Seattle, Seattle, WA, United States and **Christopher Lynnes**, NASA Goddard Space Flight Center, Greenbelt, MD, United States

**Index Terms:**

1914 Data mining [INFORMATICS]  
1918 Decision analysis [INFORMATICS]  
1926 Geospatial [INFORMATICS]  
1976 Software tools and services [INFORMATICS]

**Abstracts Submitted to this Session:**

[Sharing Planetary-Scale Data in the Cloud \(Invited\)](#) (120851)

**Joe Flasher**, Amazon Web Services, Inc, Washington, DC, United States and **Jed Sundwall**, Amazon Web Services, Global Open Data Lead, Seattle, WA, United States

See more of: [Earth and Space Science Informatics](#)



National Aeronautics and  
Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Special Thanks

## **NASA**

Mike Little  
Chris Lynnes  
Kevin Murphy

## **FSU COAPS**

Mark Bourassa  
Jocelyn Elya  
Shawn Smith  
Adam Stallard

## **NCAR**

Steve Worley  
Ji Zaihua

## **JPL NEXUS Engineers**

Kevin Gill  
Frank Greguska  
Joseph Jacob  
Nga Quach  
Brian Wilson

## **JPL Science Contributors**

Ed Armstrong  
Andrew Bingham  
Carmen Boening  
Mike Chin  
Ben Holt  
Rob Toaz  
Vardis Tsontos  
Victor Zlotnicki

## **Questions, and more information**

[Thomas.Huang@jpl.nasa.gov](mailto:Thomas.Huang@jpl.nasa.gov)