



Ecosystem Structure, Composition, and Function: Ecological Responses to Environmental Perturbations in Mammoth Mountain, California

Kerry Cawse-Nicholson, Joshua Fisher*, Caroline Famiglietti, Florian Schwandner, Ryan Pavlick, David Schimel, Amy Braverman, Robert Bogue, Jennifer Lewicki (USGS), and David Pieri

* Corresponding author




Two primary sources of uncertainty:

1) **Climate** (γ -response); versus, 2) **CO₂ fertilization** (β -response).

By far the largest sensitivity is in the β -response, with projected differences due to CO₂ fertilization in net carbon uptake up to 145% [Piao et al., 2013; Smith et al., 2016].

Plants and Elevated CO₂


- Tree responses to high CO₂ have been extensively studied in field experiments (Norby et al., 1999)
- Optimum-growth phase followed by a water-stressed phase eventually severe enough to cause death
- Threshold CO₂ concentration?



PHYSIOLOGICAL FUNCTION

TRAIT COMPOSITION


PHYSICAL STRUCTURE




Science questions

- **How do plants respond to rising CO₂**, in terms of ecosystem structure, composition, and function?
 - *Structure*: Does biomass increase with elevated CO₂?
 - *Composition*: Do canopy trait shifts occur in changing CO₂ environments?
 - *Function*: Does photosynthetic efficiency increase with increasing CO₂?








ECOLOGY

VOLCANOLOGY


Mammoth Mountain, CA

- A 1989 earthquake swarm and associated geologic deformation have drastically altered gas discharge at the land surface
- Magmatically CO₂ escapes through faults and fissures on the flanks of the mountain
- Soil CO₂ flux has created unlivable conditions for much vegetation




Mammoth Mountain, CA

- Five main tree-kill areas exist, all characterized by extreme root zone CO₂ concentrations
- Rate and spatial distribution of soil CO₂ flux: well-understood (Werner et al., Gerlach et al., Sorey et al.)
- Abundance of airborne data available in the region
- Gradient-like nature of response by Mammoth vegetation as a case study for ecological responses to future environmental perturbations, especially in vulnerable regions



Airborne

- Schwandner et al. (2017) demonstrated the potential for spaceborne detection of individual CO₂ plumes using OCO-2.
- However, greater resolution is needed to see the effect on individual trees.



Data


A wealth of information has been acquired over Mammoth mountain, including:

- **Field CO₂** measurements
- **Airborne spectrometer data**, from which we can derive various indicators of vegetation health, type, and traits
- **Thermal infrared imagery**, from which we can derive land surface temperature and evapotranspiration
- **Lidar data**, from which we can derive slope, aspect, elevation, and vegetation height
- **Fluorescence data**




AVIRIS

- AVIRIS Classic imagery was acquired in October 2014.
- L2 reflectance data was used.
- Spatial resolution is 13m.
- This scene was chosen to minimize snow cover.




MASTER

- Two MASTER datasets were obtained: one in 2013 and one in 2017.
- From the MASTER data, we derived:
 - Emissivity
 - Land surface temperature
 - Evapotranspiration
- Spatial resolution is 35m.




ASO LiDAR

- The Airborne Snow Observatory (ASO) carries out regular LiDAR surveys over Mammoth Mountain.
- This dataset was acquired in 2017.




Field CO₂

- Field CO₂ measurements were made by Werner et al. in 2013, during a larger project that has been ongoing since 1991.
- Five degassing and tree-kill areas were measured using a LI-COR® infrared gas analyzer.
- Data have been Kriged to a 1m grid.



Carbon dioxide measurements



Dataset summary

Dataset	Dates	Product	Derived Products	Resolution
AVIRIS	Oct 2014	Reflectance	Spectral features Foliar traits*	13m
MASTER	Nov 2013 Jun 2017	Thermal Reflectance	Land Surface Temperature Evapotranspiration	35m
LiDAR	Jun 2017	Point cloud	Elevation Slope, Aspect	*
Fluorescence	2017	Chlorophyll fluorescence	-	*
Carbon dioxide	Jun-Oct 2013	LiCOR soil CO2 flux	-	1m


* Data still being processed.

Hypothesis Testing Framework

- Covariate Balancing Propensity Scores stratify the data and control for confounding variables, such as rain, air temperature, and downwelling shortwave radiation.
- Exploratory data analysis on the strata, by evaluating the ability of CO₂ to predict dependent variables within each strata (i.e. removing the confounding effect) individually, in pairs, etc.

*Ma, P., Kang, E.L. Spatial Statistical Downscaling for Constructing High-Resolution Nature Runs in Global Observing System Simulation Experiments, *Technometrics*, In Review.


CO₂ versus AVIRIS NIR



* Preliminary results; effects of confounding variables still to be addressed

p<0.01


CO₂ versus Land Surface Temperature



* Preliminary results; effects of confounding variables still to be addressed

$p < 0.01$

CO₂ versus Evapotranspiration



* Preliminary results; effects of confounding variables still to be addressed

p<0.01

CO₂ versus remotely sensed ecosystem properties (total data cube)

	R ² (all data)	R ² (NDVI>0.1)	R ² (NDVI>0.3)	% remaining at 0.3
Horseshoe Lake	0.70	0.70	0.59	42%
South Side Fumarole	0.67	0.67	0.95	16%
Mammoth Fumarole	0.81	0.83	0.84	5%
Chair 12	0.55	0.57	0.57	56%
Reds Creek	0.57	0.66	0.73	16%

* Non-vegetation areas were masked.

Conclusions

- There is a **strong relationship** between CO₂ flux and our remote sensing datasets
 - R² high across all 5 ground data locations
 - Trends: as CO₂ flux increases, land surface temperature decreases and evapotranspiration increases
- Our results will have strong implications for ecological responses to elevated CO₂ in **other parts of the world**
 - This extension illustrates the need for consistent airborne data retrieval in regions where we have the most to learn, like the tropics

Ongoing work

- Once lidar data have been processed, we will use them to remove the confounding effects of slope, aspect, and elevation.
- A concurrent study is investigating the correlation between modelled ground CO₂ and parameters that may be remotely sensed.

