

IO'S NON-SO₂ SPECTRAL FEATURES

W. D. Smythe(1) L. A. Soderblom(2) and R. M. Nelson(1),(1)Jet Propulsion Laboratory/California Institute of Technology, M/S 183-601, 4800 Oak Grove Drive, Pasadena, CA 91109-8099, , (2)USGS Flagstaff, 2255 N. Gemini Drive, Flagstaff, AZ 86001-1695,

wsmythe@lively.jpl.nasa.gov, FAX 818-393-3218.

Introduction: Several absorptions have been identified in the Galileo NIMS spectra of Io that are not related to SO₂. [1,2]. These absorptions have band centers at 2.97, 3.15, 3.85, and 3.91 microns. There are also broad absorptions in the regions 1-1.3 and 3-3.4 microns. "Patterning" noise in wavelength registration, arising from the pushbroom imaging and grating motion of the NIMS instrument have previously inhibited reliable mapping of weak absorptions. Recent improvements in techniques to remove the coherent pattern noise from the NIMS dataset have been made by Soderblom. This greatly improves the signal to noise ratio and enables mapping of weak spectral signatures such as the 3.15 micron absorption on Io.

The 3.15 μ feature: The 3.15 μ feature is a very weak band having an apparent band depth of less than 4% of the continuum – in contrast to many of the SO₂ features, which have apparent band depths ranging from 15-70%. The 3.15 μ feature is reasonably ubiquitous on the surface of Io so it was seen in global average spectra obtained early in the Galileo mission [2]. The feature has not been identified but has been attributed variously to H₂S, H₂O, or OH stretch [2,3]. A map of this feature shows a variety of correlations. The feature is strongly present at lower latitudes, which suggests that it may be associated with a material less volatile than SO₂. However the absorption does appear above 30 degrees (north and south), suggesting its presence is not strictly a thermal issue. It is interesting that the absorption moves north of the equator above Pele apparently avoiding Pele's large plume deposit, is not present at Culann, yet covers Prometheus completely. The distribution at Prometheus suggests that the material leading to the 3.15 μ absorption may be associated with the Prometheus SO₂ reservoir – but it is difficult to reconcile that with the behavior of the absorption about Pele and Culann. There is a reasonably good correlation with visibly white material, but the absorption is also present where the white material is absent.

Conclusion: The 3.15 micron absorption appears to be associated with the equatorial region and with white surface features, though there are exceptions in both cases. The absorption, which is not present at Pele or Culann and appears to cover Prometheus, is probably not a direct result of plume activity.

References: [1] Shirley J. H. (2001) *BAAS* 33, #3, p1063 [2], Carlson R. W. *et al* (1997) *GRL* 24, p2479-2482, [3] Salama, F. *et al.*, (1990) *Icarus* 83 p66-82